





Biologia centrali-americana

Otto Stoll



Digitized by Google



Digitized by Google

94
108
18615

B I O L O G I A
CENTRALI-AMERICANA.

A R A C H N I D A A C A R I D E A.

BY

PROF. OTTO STOLL, M.D.

1886-1893.

Digitized by Google

CONTENTS.

	Page
INTRODUCTION	v
EXPLANATION OF THE PLATES	xvii
TROMBIDIDÆ	1, 44
ACTINIDIDÆ	6, 45
TETRANYCHIDÆ	8, 46
HYDRACHNIDÆ	9, 46
BDELLIDÆ	15, 48
EUPODIDÆ	17, 48
LEODIDÆ	18, 48
ORIBATIDÆ	24, 50
NICOLETIELLIDÆ	27, 50
GAMASIDÆ	28, 51
SARCOPTIDÆ	39, 52
INDEX	54
PLATES.	

INTRODUCTION.

COMPARED with the work that has been done in Europe upon the Arachnida Acaridea from the time of the classical writers, Linnæus, De Geer, Hermann, and Latreille, up to that of Dugès, Koch, Nicolet, P. Kramer, Mégnin, Michael, Neuman, Berlese, Canestrini, Nalepa, and many others, the American literature of this group of animals is exceedingly scanty. Thomas Say (1821), one of the first entomologists in the United States, paid some attention to this neglected group. Later on (1836), Dana and Whelpley, as well as Haldeman (1842), described and figured some North-American species of Hydrachnidæ. In our times we meet with the well-known names of Riley and Packard in connection with North-American Acarids of various groups, and Mr. Harry Garman has published a paper on the Phytoptidæ. In 1886, Messrs. Herbert Osborn and Lucien M. Underwood gave a "Preliminary List of the Species of Acarina of North America" in the 'Canadian Entomologist.' In Mexico, M. Alfred Dugès has published several valuable articles on various species of Acarids inhabiting that country, and M. Conil in Buenos Ayres has done the same for some species of the Argentine Republic.

A certain number of American Acaridea have been described by European authors, and some of the larger and more conspicuous species were already known to the founders of Entomology, to Linnæus, De Geer, and Fabricius. In modern times C. L. Koch has described several Ixodidæ from various parts of America, and Trouessart and Mégnin have studied some forms of Dermaleichidæ which live on American birds. To Antonio Berlese and R. Canestrini we are indebted for some most valuable papers on Acaridea from the La Plata region and from Brazil.

But nevertheless we are far from possessing a knowledge of the American Acarid fauna comparable to that which we have long had of the European forms.

In the following pages an attempt has been made to fill, to some extent, the gap which at present separates the forms known from North America and those described by

the above-quoted Italian authors from various parts of South America, and to characterize some new types from Mexico and Central America. The materials which I had at my disposal were derived from various sources. During a stay of nearly five years in various parts of Guatemala, I tried to make myself acquainted with the Acarid-fauna of that country, by studying, as far as the unsettled life of a medical practitioner would allow, the living forms. I had then with me a Hartnack microscope of but moderate powers (objectives 4 and 7), the lenses of which became damaged by the long influence of the excessive humidity of the climate of the Costa Grande. Moreover, I laboured under an almost total want of modern literature on Acarida, having only some of the works of the older writers with me. These unfavourable circumstances will account to some extent for the differences the reader may find in the execution of the drawings and the descriptions of some species, several of which are too delicate for preservation in alcohol, the only method then within my reach. Many of my drawings, especially of Gamasidae, have been entirely omitted from the present memoir, as being too incomplete to allow a comparison with the European forms; and even amongst those which I have admitted there are some which I should have liked to revise again from the specimens themselves.

Fortunately, this lack of preserved materials has in numerous cases been made up by dried specimens obtained by other naturalists who have travelled in Central America.

The fact that other collectors have frequently fallen in with the same species as myself shows that my researches, though far from complete, were sufficient at least to give an idea of the composition of the Acarid fauna of a tropical country, and to enable me to come to some general conclusions as to the geographical distribution of the various groups.

The most striking fact elicited by the study of the Acarid fauna of Central America is undoubtedly the great similarity between the types with which we are familiar in the temperate regions north of the European Alps and those occurring in the gloomy shadow of the tropical forest. Under the bark and in the fissures of putrefying tree-trunks in the tropics we meet with some minute Gamasidae which only by a close microscopical examination can be distinguished from European species living under similar conditions. Upon various beetles of the families Copridae and Passalidae are found, with others, such well-known forms as *Gamasus (Holostaspis) marginatus* and *G. coleopterorum*, species already described by authors of the past century. Among the fallen leaves and in the decaying fruits of *Theobroma*, *Lucuma*, and other tropical

trees, which lie scattered over the damp ground of the forest, we frequently observe the slow-moving Oribatidæ, or the soft velvety Trombididæ, the larvæ of which may be found attached to the wings of dragon-flies, grasshoppers, and other insects. The larger species of Muscidæ carry on their thorax the hypopial forms of Tyroglyphidæ, just as they do in Europe; and the Belostomidæ, which we find in the ponds of the high valleys and barrancas of Guatemala, have their legs infested by the larvæ of Hydrachnidæ, in the same way as are those of *Ranatra* and *Nepa* in Europe. It is a remarkable circumstance that not one of the species described in the following memoir represents a generic type entirely new or peculiar to Central America. With the exception of the holotropical genus *Megiathanus* (fam. Gamasidæ), and the chiefly American genus *Amblyomma* (fam. Ixodidæ), all the genera enumerated by me likewise occur in the Palearctic Region.

Geographical Distribution of the Genera of Acarids hitherto found in Central America.

Fam. TROMBIDIDÆ.

Gen. **TROMBIDIUM.**

- Palearctic Region (Europe, N.E. Siberia).
- Nearctic R. (U.S.A.).
- Neotropical R. (Central and South America).
- Æthiopian R. (Senegambian, Gold Coast, S.W. Africa, Cape of Good Hope).

Gen. **RYNCHOLOPHUS.**

- Palearctic Region (Europe).
- Nearctic R. (U.S.A.).
- Neotropical R. (Central and South America).

Gen. **LINOPODES.**

- Palearctic Region (Europe).
- Neotropical R. (Guatemala, Buenos Ayres).

Fam. ACTINEDIDÆ.

Gen. **ACTINEDA.**

- Palearctic Region (Europe, Siberia).
- Neotropical R. (Central and South America).

Fam. TETRANYCHIDÆ.

Gen. **TETRANYCHUS.**

- Palearctic Region (Europe).
- Nearctic R. (U.S.A.).
- Neotropical R. (Mexico, Guatemala).

Fam. HYDRACHNIDÆ.**Gen. ATAX.**

- Palearctic Region (Europe).
- Nearctic R. (U.S.A.).
- Neotropical R. (Mexico †, Guatemala).
- Æthiopian R. (East Africa ‡).

Gen. NEREA (*Curipes*).

- Palearctic Region (Europe, Bering's Island).
- Neotropical R. (Mexico, Guatemala).
- Æthiopian R. : East Africa ‡.

Gen. LIMNÆIA.

- Palearctic R. (Europe).
- Nearctic R. (U.S.A.).
- Neotropical R. (Guatemala).
- Æthiopian R. (German East Africa ‡).

Fam. BDELLIDÆ.**Gen. BDELLA.**

- Palearctic Region (Europe, N.E. Siberia, Bering's Island).
- Nearctic R. (U.S.A.).
- Neotropical R. (Guatemala, Brazil, Paraguay, La Plata).

Fam. EUPODIDÆ.**Gen. SCYPHICÆ.**

- Palearctic Region (Europe, N.E. Siberia, Japan).
- Neotropical R. (Guatemala).

Fam. IXODIDÆ.**Gen. IXODES.**

- Palearctic Region (Europe, Bering's Island).
- Nearctic R. (U.S.A.).
- Neotropical R. (Central and South America).
- Æthiopian R. (South Africa [*teste* C. L. Koch]).

Gen. AMBLYOMMA.

- Nearctic Region (U.S.A.).
- Neotropical R. (Central and South America).
- Oriental R. (Java, Philippine Islands, Butang, Continental India [*teste* C. L. Koch]).

* The Mexican species described by M. Alfred Dugès as *Atax alzatei* belongs, according to a private communication of Herr F. Koenike, to the genus *Nesoa*.

† Herr F. Koenike, the well-known specialist in this group, writes me that he possesses specimens of *Atax* collected by Dr. Stuhlmann during his expedition with Emin Pasha.

‡ According to the above-quoted communication of Herr F. Koenike.

INTRODUCTION.

ix

Australian R. (New Holland [*teste* C. L. Koch]).

Æthiopian R. (South Africa [*teste* C. L. Koch]).

Gen. ARGAS.

Palearctic Region (Europe, Egypt, Persia).

Nearctic R. (U.S.A.).

Neotropical R. (Mexico, Guatemala, Paraguay).

Æthiopian R. (Ovampo Land).

Fam. ORIBATIDÆ.

Gen. ORIBATA.

Palearctic Region (Europe, N.E. Siberia, Algeria).

Nearctic R. (U.S.A.).

Neotropical R. (Central America, Brazil, Paraguay).

Gen. HOLOPHORA.

Palearctic Region (Europe).

Nearctic R. (U.S.A.).

Neotropical R. (Guatemala, Brazil).

Fam. NICOLETIELLIDÆ.

Gen. NICOLETIELLA.

Palearctic Region (Europe).

Neotropical R. (Guatemala).

Fam. GAMASIDÆ.

Gen. UROGOMA.

Palearctic Region (Europe).

Nearctic R. (U.S.A.).

Neotropical R. (Central America, Brazil, Paraguay).

Gen. HOLOSTAPHIS.

Palearctic Region (Europe, Bering's Island).

Neotropical R. (Central America, Brazil, Paraguay, La Plata).

Gen. GAMASUS.

Palearctic Region (Europe, Japan).

Nearctic R. (U.S.A.).

Neotropical R. (Central America, Brazil, Paraguay, La Plata).

Gen. CELENOPSIS.

Palearctic Region (Europe).

Neotropical R. (British Honduras, Panama, Brazil, Paraguay, La Plata).

Gen. PACHYLELAPH.

Palearctic Region (Europe).

Neotropical R. (Mexico, Brazil, Paraguay).

BIOL. CENTR.-AMER., Arachu, Acar., January 1893.

b

Gen. *MESOSTRANCUS*.

- Oriental Region (Malacca).
- Australian R. (New Guinea, Queensland).
- Neotropical R. (Mexico, Guatemala, Paraguay).
- Æthiopian R. (Gold Coast *).

Gen. *DERMANTSEUS*.

- Palaearctic Region (Europe).
- Neartic R. (U.S.A.).
- Neotropical R. (Guatemala).

Fam. *SARCOPTIDÆ*.Gen. *TYROGLYPHUS*.

- Palaearctic Region (Europe).
- Neartic R. (U.S.A.).
- Neotropical R. (Guatemala).

Gen. *MESONINIA*.

- Palaearctic Region (Europe).
- Neotropical R. (Guatemala).

Gen. *PREBOLICUS*.

- Palaearctic Region (Europe).
- Neartic R. (U.S.A.).
- Neotropical R. (Guatemala, Brazil, Guiana, Ecuador, Patagonia).
- Oriental R. (Himalaya, India, Java, Philippine Islands, China).
- Australian R. (New Holland, New Guinea, New Zealand, New Caledonia, Tahiti, &c.).
- Æthiopian R. (Gold Coast and other parts of Africa).

Gen. *PROCTOPHYLLODES*.

- Palaearctic Region (Europe).
- Neotropical R. (Guatemala).

There can be no doubt that the geographical distribution of the various families, genera, and even species of many Acarids will eventually be found to be much larger. Several are already known to extend over more than one of the zoo-geographical regions, *i. e.* :—

Atax crassipes, O. F. Müll. †—Europe, Guatemala.

Actinoda baccharum, L. ‡—Europe, Guatemala, Paraguay (Rio Apa), Buenos Ayres.

Holostaspis marginatus, Herm.—Europe, Guatemala, Brazil (Matto Grosso), Paraguay (Asuncion, Rio Apa), La Plata (Buenos Ayres).

Undoubtedly the migrations of the Acarids may account, to some extent, for the uniformity of types, especially as regards the wholly or partially parasitical species.

* I possess a new and very interesting species belonging to this remarkable genus from Acera (Gold Coast), which I propose shortly to describe elsewhere.

† See page 47.

‡ See page 45.

In some instances, as in the *Dermaleichidæ*, the passive migration on birds' wings is no doubt the usual mode of dispersion; and as the regions of Central America form a sort of rendezvous and winter station for many birds of passage which are infested by *Dermaleichidæ*, contact and interchange of Acarid forms of this particular group are continually taking place. Much less extensive, though yet considerable, is probably the passive migration and the dispersion of those Acarids which, in some stages of their existence, adhere to insects and to bats, and which, therefore, are carried by insects or on bats' wings, such as various Tyroglyphidæ, Gamasidæ, Trombididæ, and Hydrachnidæ; the Hydrachnidæ attach themselves to Hemipterous water-insects which at night abandon their ponds and take wing. In the course of many successive generations these Acarids may thus spread over large areas. Amongst the non-parasitic species, the active migration of those of open habits and rapid locomotion, such as the Trombididæ, Actinedidæ, &c., may influence the dispersion of types and the mutual penetration into different faunas. How far the aerial transportation by wind and storm, which are such powerful agencies in the migration of winged insects and young spiders, may facilitate the passive migration of Acarids is, as yet, entirely unknown.

But, taken as a whole, the various modes of migration, numerous as they are, only serve to explain the similarity of types in more or less contiguous land-areas, such as North, Central, and South America, and the wide range of single species. The almost universal occurrence, however, of certain genera, such as *Ixodes*, *Argas*, *Actineda*, *Trombidium*, *Rhyncholophus*, *Holostaspis*, *Oribata*, *Atax*, and the world-wide distribution of the fundamental types of Acarids, must have another and more general cause. This, most probably, is owing to the early dispersion of the primary Acarid types from their centres of origin, and in the comparative persistency of those types, due to a relatively perfect correspondence between the once acquired differentiation of their essential organs and their modes of life. Bearing in mind the fact that the local faunas of two so very different and widely separated regions as Central Europe and Central America possess a comparatively large number of identical generic types and of closely allied species of Acarids, we have perhaps a right to generalize and to presume:—

(1) That this uniformity is, geologically speaking, very ancient, and originated in a comparatively early geological period when the relative positions of the continents, the islands, and the seas were altogether different from what they are now.

(2) That the Acarids long ago arrived at that degree of organic development (progressive or regressive) which was the fittest for their various modes of life, not partaking in the great and rapid changes of generic and specific characters which, in the course of the more recent geological epochs, have more or less affected so many of the higher organized types of the animal kingdom.

The presence of *Megisthanus* in Central America, a well-characterized Gamasid genus which has not yet been found in any of the extratropical regions, is a fact of peculiar interest. In his original paper Signor T. Thorell* described three species from Java and two from New Guinea. One of the latter (*M. testudo*, Thor.) has also been mentioned by Signor G. Canestrini† as occurring in Queensland. One species has since been added by Signor A. Berlese from Paraguay‡, and most probably the "Gamasé géant" of A. Dugès§, from Brazil, belongs to the same genus. I have already had occasion to mention the fact that this remarkable type also occurs in tropical Africa||. It therefore belongs to the tropical regions of not less than four different zoo-geographical areas, viz.: the Oriental (Java), the Australian (New Guinea and Australia), the Neotropical (Central and South America), and the Ethiopian (Gold Coast)—a fact which it would be very difficult to explain by a migratory dispersion of recent origin from one starting-point. It is far more reasonable to regard these now so widely dispersed *Megisthani* as the surviving members of a once, that is in former geological periods, coherent group of Gamasids which have been separated in consequence of the slow but material changes of the earth's surface, principally by the successive breaking down of large masses of the earth's crust and the filling up of the thus formed gulfs by the seas.

The genus *Megisthanus* is by no means the only example of the occurrence of one and the same animal type at different regions which at the present time are separated by large tracts of sea, and which for long periods have not had any direct land communication whatever with one another.

* T. Thorell, Descrizione di alcuni Aracnidi inferiori dell' Arcipelago Malese, 1882.

† G. Canestrini, Acari nuovi o poco noti, p. 14 (Atti del R. Istituto Veneto di Scienze, Lettere ed Arti, t. ii. ser. vi. 1884).

‡ A. Berlese, Acari Austro-Americani, 1888.

§ A. Dugès, Recherches sur l'ordre des Acariens, 3^e Mém., 1837.

|| See above, p. x, note.

The remarkable distribution of some of the higher animals, such as the *Prosimia* among mammals, the *Ratitæ* among birds, the *Crocodylida* among reptiles, is well known, and every student of terrestrial Invertebrata who has paid some attention to the geographical distribution of his favourite group must be acquainted with similar facts. I may be allowed here to mention the Gasteropod genus *Clausilia*, the *Nenia* group of which is now limited to the high valleys and mountain chains of Peru, Ecuador, and Colombia, and, with one species only, to the island of Puerto Rico, and which has its nearest allies not in the New, but in the Old World, in the *Lamifera* group (*Neniatlanta*, Bgt.), which now lives on the top of La Rhune, a mountain near the coast of the Bay of Biscay, and in the *Garnieria* group, the members of which inhabit the mountainous districts of China, Siam, and Cambodia. On some pieces of bark in the virgin forests of the Pacific slope of Guatemala I discovered a new species of *Diplommatina**, a Gasteropod type, the autochthonous members of which had previously only been known from India and the neighboring archipelagos. In the woods near Retalhuleu (N.W. Guatemala) I found a new species of the Myriopod genus *Polyxenus*, the type of which is the well-known *P. lagurus*, L. Another species of this well-defined Chilognath genus has been mentioned by Mr. Humbert from Ceylon, and one species has been described from North America by Say. A not less characteristic Myriopod type, the genus *Siphonophora*, abounds under the bark of the fir trees near the summits of the volcanos Agua and Fuego in Guatemala, whilst a nearly allied species has been found in Madagascar by my friend, Prof. C. Keller. Another species has been described from Ceylon by Mr. Humbert. In the woods of the Volcan de Agua, at an elevation of about 10,000 feet, I met with a species of land-leech belonging to the genus *Cycticobdella*, Grube, which is closely allied to, if not identical with, *C. lumbricoides*, Gr., discovered by Prof. Frits Müller at Desterro in Brazil.

Similar instances of an almost world-wide distribution might, no doubt, be found among other groups of Invertebrata whose facilities for active or even passive migration are very limited.

Unfortunately our present knowledge of the Acarids is too fragmentary to allow any more definite speculations as to the phylogeny of this group. Whether it is, geologically speaking, as ancient as some other groups of the Arachnida, or

* This species has since been described and figured as *Diplommatina stelli* in the 'Biologia Centrali-Americana' (Mollusca, p. 20, Tab. I. figg. 19 a, b), by Prof. E. von Martens.

whether it is of a more recent origin, whether its various families, some of which are not very closely allied to each other, took their origin from one or from several types, are questions we shall probably never be able to answer. Arachnoid Arthropods appear early in the strata of the primary periods, and it is quite possible that Acaroid types were among them, though the delicacy and minuteness of their structure made their preservation highly improbable. It is even possible that a closer examination of those sedimentary layers which are fine enough for the preservation of more delicate organisms, such as some of the tertiary strata (those of Oeningen for instance), may lead to the discovery of the larger and more chitinized forms, such as the Ixodidæ, Gamasidæ, and Oribatidæ. At present only one Acarid species is known from the Tertiary deposits of Oeningen. A larger number of Acarid types have been described from the Baltic amber. The brown coal of Rott and the Green River beds of Wyoming have furnished a few isolated forms.

But though, as yet, any direct proofs of the geological antiquity of the Acarid type beyond the Oligocene are wanting, the above-quoted fact of a most extensive geographical distribution of the principal genera, and the general uniformity and similarity of the European and extra-European local faunas, as far as they are known at present, are highly in favour of a pretertiary origin of the Acarid types.

If we compare the Acarid fauna of those parts of Europe where it has been somewhat carefully studied with that of Central America, we are compelled to say that Central America is comparatively poor as regards the number of species, far more so than we should be inclined to anticipate when we take into consideration its great variety of soil and climate and its general richness in vegetable and animal productions. I willingly admit that the districts to which my personal researches were confined are not very extensive, and perhaps other parts of Guatemala, such as the high valleys of Alta Vera Paz, or the forests of the alpine mountains of the "Altos," or even the richer slopes of the Atlantic coast, may have a more varied Acarid fauna; but as the various collectors in other parts of Central America did not, in so conspicuous a group as the Oribatidæ, meet with any other species than those which I obtained in Guatemala, it seems to me probable that even those parts, when searched more carefully, will not prove very much richer or more varied in Acarid forms, though undoubtedly they may yield some new species which escaped my notice. I am therefore inclined to believe that there really exists a comparative scarcity of Acarid species, at least in Western Guatemala, and that the principal cause of this is to be found in

the atmospheric and climatic conditions of that country. For organisms of such delicate construction, the hygrometric equilibrium of which is so easily disturbed, the extreme aridity of the "verano" or dry season, which in Western Guatemala lasts for several months, must prove much more fatal than does even the European winter with its frozen and snow-covered ground. On the other hand, the torrential "agnaceros" of the "invierno" or rainy season probably cause the destruction of numerous non-parasitic soft-bodied and unprotected Acari. It is for these reasons probably that we find the greatest variety of species in those genera which are protected to some extent by their entirely or partially parasitic habits, such as the Ixodidae, Gamasidae, Hydrachnidae, and Dermalichidae, or which, like *Trombidium*, are capable of a somewhat rapid locomotion, which enables them to reach, in case of need, a shelter to protect them from being drowned or dried up. The various stages of the tough-skinned "Garrapatos" (ticks) are enabled, however, to withstand alike the heat and dryness of the "verano" and the deluges of the "invierno."

The limited number of species is, in some instances at least, counterbalanced to a certain extent by an abundance of individuals. This is the case in some species which are more resistant, or better protected against the influences of the climate, than the majority of their congeners—for example, with *Trombidium muricola*, *Tetranychus guatemalæ-novæ*, *Atax allicola*, *Amblyomma mixtum*, *Oribata centro-americana*, *Holostaspis marginatus*, and the various species of Dermalichidae.

I regret that I had not sufficient opportunities for observing the vertical distribution of the Central-American Acarida. My ascents of the volcanos Agua and Fuego were both made in the dry season, in the unfavourable months of January and February, when, at night, the temperature on the summits was as low as -2°C ., and when, from cold and aridity, Arthropod life was reduced to a few species of Insects, Myriopods, and Spiders, these living under the bark of the scattered fir trees and under stones*. But, judging from what I have observed in the Swiss Alps, where I found a small number of such conspicuous types as *Rhyncholophus*, *Erythraeus*, *Oribata*, and *Gamasus*

* When I passed the night in the crater of the Volcan de Agua, 19/20 February, 1881, I found in the morning the water in our jar covered with a crust of ice. Under the bark of a fir tree near the edge of the crater I found the dead body of a small species of venomous snake, which had evidently been killed by the cold. Notwithstanding, I discovered under a flat stone in the crater itself an ants' nest, the inhabitants of which were winged. This species has since been described by my friend, Prof. A. Forel, as *Leptothorax stollii*.

reaching the snow-line, and where one of them at least (*Rhyncholophus nivalis*, Heer) even surpasses it *, I am inclined to believe that within the tropics Acarids will be found at as high an elevation as any other Arthropod group.

PROF. OTTO STOLL, M.D.

Küssnacht, near Zürich, January 1893.

* According to the observation of the late Prof. Heer, *Rhyncholophus nivalis* reaches an elevation of 9580 feet (top of the Pin Levarone).

EXPLANATION OF THE PLATES.

TAB. I.

- Fig. 1. *Trombidium mexicanum* (pp. 1, 44): 1 a, palpus; 1 b, mandible; 1 c, first tarsus; 1 d, second tarsus.
2. *Trombidium trilineatum* (pp. 4, 45): 2 a, palpus; 2 b, tarsus; 2 c, texture of the skin.
3. *Trombidium albicollis* (pp. 5, 45): 3 a, palpus.

TAB. II.

- Fig. 1. *Trombidium hispidum* (pp. 2, 44): 1 a, palpus; 1 b, mandible; 1 c, first tarsus; 1 d, second tarsus.
2. *Trombidium guayanicola* * (pp. 4, 45): 2 a, palpus; 2 c, tarsus.
3. *Trombidium muricola* (pp. 5, 45): 3 a, 3 b, varieties of the same.

TAB. III.

- Fig. 1. *Trombidium narutsum* (pp. 2, 44): 1 a, palpus; 1 b, second tarsus; 1 c, first tarsus; 1 d, anterior margin of sternum; 1 e, mandible; 1 f, texture of the skin; 1 g, skin with hairs.
2. *Bdella splendida* (pp. 15, 48): 2 a, tarsus; 2 b, top of the rostrum; 2 c, mandible.
3. Larva of *Bdella* (?) sp.† (p. 16): 3 a, first tarsus, showing the texture of the skin; 3 b, mandible; 3 c, palpus; 3 d, second tarsus.

TAB. IV.

- Fig. 1. *Trombidium quinque-maculatum* (pp. 3, 45): 1 a, palpus; 1 b, fourth joint of first leg; 1 c, first tarsus.
2. *Rhyncholophus erinaceus* (pp. 6, 45): 2 a, palpus, with top of labium; 2 b, section of leg with hairs; 2 c, tarsus.

* = *T. muricola*, var.

† Probably belongs to *Rhyncholophus*.

TAB. V.

- Fig. 1. *Actinoda flavola** (pp. 7, 45): 1 *a*, ventral surface.
 2. *Actinoda antiguensis* (pp. 7, 45): 2 *a*, palpus; 2 *b*, mandible; 2 *c*, tarsus (lateral view);
 2 *d*, tarsus (dorsal view).
 3. *Actinoda retalteca* (pp. 7, 45): 3 *a*, tarsus (lateral view); 3 *b*, mandible; 3 *c*, tarsus
 (ventral view).

TAB. VI.

- Fig. 1. *Tetranychus guatemala-nova* (pp. 8, 46): 1 *a*, palpi; 1 *b*, tarsus with ambulacra;
 1 *c*, acicular mandibles.
 2. *Scyphius maniacus* (pp. 17, 48): 2 *a*, last joint of palpus; 2 *b*, mandible; 2 *c*, tarsus;
 2 *d*, hypostome.

TAB. VII.

- Fig. 1. *Atax alticola* (pp. 9, 46): 1 *a*, ventral surface; 1 *b*, dorsal surface of another specimen
 showing unusual position of eggs; 1 *c*, fourth joint of hind leg, showing the
 swimming bristles; 1 *d*, fourth joint of first leg; 1 *e*, claws; 1 *f*, palpus; 1 *g*, genital
 laminae.
 2. *Limnesia guatemalteca* (pp. 13, 46): 2 *a*, ventral surface; 2 *b*, palpus; 2 *c*, genital
 laminae; 2 *d*, tarsus of hind leg; 2 *e*, top of the mandible.
 3. *Limnesia puteorum* (pp. 14, 48): 3 *a*, palpus; 3 *b*, genital laminae; 3 *c*, tarsus of
 third leg.

TAB. VIII.

- Fig. 1. *Atax septem-maculatus*† (pp. 9, 46, 47): 1 *a*, ventral surface; 1 *b*, mandible;
 1 *c*, palpus; 1 *d*, genital laminae; 1 *e*, hair of first leg.
 2. *Limnesia lata* (pp. 14, 48): 2 *a*, part of ventral surface, showing the disposition of the
 epimera and genital laminae; 2 *b*, palpus; 2 *c*, mandibles; 2 *d*, tarsus of hind leg.

TAB. IX.

- Fig. 1. *Atax septem-maculatus*, var. *ypsilou*† (pp. 10, 47): 1 *a*, ventral surface; 1 *b*, genital
 laminae; 1 *c*, palpus.
 2. *Limnesia longipalpis* (pp. 13, 47): 2 *a*, part of the ventral surface, showing the dispo-
 sition of the epimera and genital laminae; 2 *b*, mandible; 2 *c*, palpus.

* *A. flavola*, *A. antiguensis*, and *A. retalteca* = *A. bocurum*, Lion., varr.

† Probably nymphal stage of *A. alticola*.

TAB. X.

- Fig. 1. *Atax dentipalpis*, ♀ (pp. 10, 47): 1 a, ventral surface; 1 b, palpus; 1 c, mandible; 1 d, posterior margin of the abdomen, with the genital laminae (ventral view).
 2. *Nesoa guatemalensis*, ♀ (pp. 11, 47): 2 a, part of the ventral surface, showing the disposition of the epimera and genital area; 2 b, palpus.

TAB. XI.

- Fig. 1. *Nesoa guatemalensis*, ♂ (pp. 11, 47): 1 a, part of the ventral surface, showing the disposition of the epimera and genital laminae; 1 b, genital laminae; 1 c, palpus; 1 d, mandible; 1 e, section of the fourth joint of hind leg; 1 f, tarsus of first leg.
 2. *Nesoa nuxulus* (pp. 12, 47): 2 a, part of the ventral surface, showing the disposition of epimera and genital area; 2 b, genital area; 2 c, palpus.

TAB. XII.

- Fig. 1. *Amblyomma mixtum*, ♀ (pp. 19, 49): 1 a, the same, full of blood after suction; 1 b, genital aperture; 1 c, anal aperture; 1 d, top of mandible (pseudochela); 1 e, stigmatic plate (peritreme); 1 f, radula of the area maxillaris, viewed from beneath; 1 g, tarsus of hind leg; 1 h, palpus; 1 i, tarsus of first leg.
 2. *Amblyomma mixtum*, ♂ (pp. 19, 49): 2 a, genital aperture; 2 b, anal aperture.
 3. *Amblyomma foreli*, ♀ (pp. 21, 50), tarsus of hind leg: 3 a, top of mandible (pseudochela); 3 b, palpus of the same. (See also Tab. XIV, fig. 3.)

TAB. XIII.

- Fig. 1. *Ixodes boarum*, ♀ (pp. 18, 19): 1 a, ventral surface; 1 b, dorsal surface of a young individual; 1 c, sucking-apparatus, showing the palpi, the radula of the area maxillaris, and the mandibles (pseudochela); 1 d, tarsus; 1 e, stigmatic plate (peritreme). (See also Tab. XIV, fig. 4.)

TAB. XIV.

- Fig. 1. *Amblyomma crassipunctatum*, ♂ (pp. 22, 50): 1 a, anal aperture; 1 b, genital aperture; 1 c, stigmatic plate (peritreme); 1 d, palpus; 1 e, tarsus of first leg; 1 f, tarsus of hind leg; 1 g, top of the mandible; 1 h, radula of the area maxillaris.
 2. *Amblyomma sabanerae*, ♀ (pp. 23, 50): 2 a, coxa of hind leg; 2 b, tarsus of hind leg; 2 c, tarsus of first leg; 2 d, radula of the area maxillaris; 2 e, stigmatic plate (peritreme); 2 f, palpus; 2 g, top of the mandible; 2 h, genital aperture; 2 i, anal aperture.
 3. *Amblyomma foreli*, ♀ (pp. 21, 50): 3 a, stigmatic plate (peritreme); 3 b, first tarsus; 3 c, genital aperture; 3 d, anal aperture. (See also Tab. XII, figs. 3-5.)
 4. *Ixodes boarum*, ♀ (pp. 18, 49), anal aperture. (See Tab. XIII.)

TAB. XV.

- Fig. 1. *Oribata centro-americana* (pp. 24, 50): 1 *a*, ventral surface; 1 *b*, lateral view of the cephalothorax with pteromorphæ; 1 *c*, mandible; 1 *d*, palpus with part of the epistomium; 1 *e*, tarsus; 1 *f*, eggs.
2. *Oribata regifrons* (pp. 25, 50): 2 *a*, ventral surface; 2 *b*, mandible; 2 *c*, leg; 2 *d*, palpus.
3. Larva of *Oribata* sp. (p. 26): 3 *a*, mandible; 3 *b*, palpus with part of the epistomium; 3 *c*, tarsus; 3 *d*, hair.
4. *Hoplophora retalteca* (ventral view) (pp. 27, 50): 4 *a*, lateral view; 4 *b*, mandible; 4 *c*, epistomium, with right palpus of the retracted mandibles; 4 *d*, palpus; 4 *e*, tarsus; 4 *f*, eggs.

TAB. XVI.

- Fig. 1. *Nicolettiella neotropica* (pp. 27, 50): 1 *a*, palpi; 1 *b*, microscopical texture of the epiderm; 1 *c*, mandible.
2. *Uropoda echinata* (pp. 28, 51): 2 *a*, hypostome and palpi; 2 *b*, tarsus; 2 *c*, microscopical texture of the epiderm; 2 *d*, chela of the mandible; 2 *e*, tarsus of first leg.
3. *Uropoda inaequipunctata* (pp. 29, 50): 3 *a*, ventral view; 3 *b*, chela of the mandible; 3 *c*, tarsus; 3 *d*, anal orifice.
4. *Celanopsis uropodoides*, ♀ (pp. 35, 52), ventral view: 4 *a*, chela of the mandible; 4 *b*, tarsus of first leg; 4 *c*, genital orifice; 4 *d*, microscopical texture of the skin. (See also Tab. XIX. figs. 3-3 *b*.)

TAB. XVII.

- Fig. 1. *Uropoda centro-americana* (pp. 30, 51): 1 *a*, buccal parts in the ventral view, coxæ of first leg, palpi, corniculi labiales, and labium—the whole protected by the anterior margin of the dorsal plate (the mandibles are wanting); 1 *b*, tarsus of first leg; 1 *c*, tarsus of second leg; 1 *d*, chela of the mandible; 1 *e*, sternal plate; 1 *f*, spiraculum.
2. Nymph of *Uropoda centro-americana* (pp. 30, 51): 2 *a*, ventral view; 2 *b*, tarsus.
3. *Uropoda piriformis* (pp. 31, 51): 3 *a*, ventral view, without the legs; 3 *b*, tarsus; 3 *c*, hypostome and palpi; 3 *d*, chela of the mandible.
4. *Uropoda discus* (pp. 29, 51): 4 *a*, ventral view of the first pair of legs, palpi, and mandibles; 4 *b*, palpus; 4 *c*, tarsus of second leg.

TAB. XVIII.

- Fig. 1. *Megisthanus gigantodes*, ♀ (pp. 31, 51): 1 *a*, ventral view; 1 *b*, genital orifice; 1 *c*, tarsus of second leg; 1 *d*, ventral view of the buccal parts.
2. *Megisthanus gigantodes*, ♂ (pp. 31, 51): 2 *a*, ventral view; 2 *b*, chela of the mandible (deprived of its appendages by maceration in a solution of caustic potash); 2 *c*, ventral

EXPLANATION OF THE PLATES.

xxi

view of the buccal parts; 2 *d*, genital orifice; 2 *e*, microscopical texture of the epiderm; 2 *f*, adhesive grooves at the end of the sternal plate; 2 *g*, chela of the mandible with its appendages.

TAB. XIX.

- Fig. 1. *Megisthanus armiger*, ♂ (pp. 34, 51): 1 *a*, ventral view; 1 *b*, lip and palpi; 1 *c*, chela of the mandible; 1 *d*, genital orifice; 1 *e*, tooth-bearing joints of second leg.
2. *Pachylolaps haeros*, var. *mexicanus*, ♂ (pp. 37, 38, 52): 2 *a*, ventral view; 2 *b*, chela of the mandible; 2 *c*, genital orifice; 2 *d*, ventral view of the buccal parts, lip, corniculi labiales, and palpi; 2 *e*, leg of the second pair.
3. *Celanopsis uropodoides*, ♀ (pp. 35, 52), dorsal view: 3 *a*, chela of the mandible with its appendages; 3 *b*, lip with corniculi labiales. (See also Tab. XVI. figg. 4-4*d*.)
4. *Celanopsis megisthanoides*, ♀ (pp. 36, 52): ventral view of the buccal parts, with the lip, corniculi labiales, and palpi.

TAB. XX.

- Fig. 1. *Celanopsis megisthanoides*, ♀ (pp. 36, 52): 1 *a*, ventral view; 1 *b*, chela of the mandible.
2. *Holostaspis marginatus* (pp. 39, 52): 2 *a*, ventral view; 2 *b*, lip, corniculi labiales, and palpi; 2 *c*, chela of the mandible; 2 *d*, tarsus of second leg.

TAB. XXI.

- Fig. 1. *Pterolichus momotorum*, ♂ (pp. 39, 52): 1 *a*, anal appendages with the adhesive discs; 1 *b*, chela of the mandibles and palpi.
2. *Pterolichus momotorum*, ♀ (pp. 39, 52): 2 *a*, ventral view.
3. *Proctophylodes sialiarum*, ♂ (pp. 42, 53).
4. *Proctophylodes sialiarum*, ♀ (pp. 42, 53): 4 *a*, leg of the second pair; 4 *b*, chela of the mandible; 4 *c*, egg with embryo.
5. *Megninia pteroglossorum*, ♂ (pp. 40, 52): 5 *a*, nymph of ♀; 5 *b*, larva.

BIOLOGIA CENTRALI-AMERICANA.

ZOOLOGIA.

Class ARACHNIDA.

Order ACARIDEA.

Suborder I. ACARINA-TRACHEATA, Kramer.

Fam. TROMBIDIDÆ.

[Kramer, Grundsätze zur Systematik der Milben, in Arch. für Naturg. xliii. p. 226 (1877).]

TROMBIDIUM.

Trombidium, Latreille, Gen. Crust. et Ins. i. pp. 144, 145 (1806).

1. *Trombidium mexicanum*, sp. n. (Tab. I. figg. 1-1 d.)

Corpus oblongum, pyriforme, depressum, postice rotundatum, tomentoso-hirsutissimum; tomentum e pilis quadrifidis sistens, colore uniformi coccineo, ex sericeo nitente; superficies dorsalis impressionibus transversalibus induta. Pedes breviusculi, coccineo-sericei, tomentosi; subtus (exceptis tarsis) pilis longis pinnatis in seriem dispositis instructi. Palpi longi, marginem corporis anteriorum superantes; pilis tomentosis pinnatis ladati; appendicula lobiformi lanceolata, basid lata, longe pilosa. Mandibulae fusiformes, ungue falciformi acuto armatae. Oculi coccinei, in tomento fere absconditi.

Long. 5-6, lat. max. 4 millim.

Hab. MEXICO, Presidio (*Forrer*). Two examples.

Body oblong, pyriform, depressed from above and below; shoulders protracted between the coxae of the second and third pairs of legs; posterior part of the body cylindrical, its hind margin rounded; dorsal surface with a few transverse impressions; colour uniform, a bright scarlet; the whole body thickly covered with velvety, quadrifid hairs. Coxae of the second and third pairs of legs separated by the protracted shoulders. Legs rather short, when compared to the mass of the body, of about equal length, the anterior ones a trifle thinner and longer than the others; slightly compressed, higher than broad, thickly covered with short red hairs which give them a whitish silky lustre; beneath bearing brushes of long, straight, pinnate hairs, which in the fore legs are but imperfectly developed and limited to the first three joints (counting from the coxa); the brushes are wanting from all the tarsi; front tarsi obliquely truncate at the top, their lower surface covered with very short hairs which are arranged into a sort of tactile brush, their claws much smaller than those

BOL. CENTR.-AMÉR., Arachn. Acar., December 1886.

B*

of the other pairs; the tarsi of the second, third, and fourth pairs bearing long pinnate hairs. Palpi long, extending far beyond the front margin of the body; covered with velvety, pinnate hairs, which on the surface of the appendicula form long fringes. The fusiform mandibles are terminated by a broad, falciform, sharply pointed claw, which is much shorter than the mandible itself.

The description of this beautiful species is drawn from two dried and pinned specimens, which were collected by Mr. Forrer.

2. *Trombidium hispidum*, sp. n. (Tab. II. figg. 1-1 d.)

Corpus ovato-trigonum, margine anteriori in apicem trigonum protracto; immaculatum, tomentoso-hirsutum, uniformiter coccineum; pilis rarioribus, disseminatis, elongatis, pinnatis, ex tomento exsertis indutum, precipue in regione humerali et versus marginem posteriorem. Pedes inaequales, in paribus 1^o et 4^o longiores; tarsus primi paris fusiformis, unguibus minimis. Palpi exserti, pilis elongatis induti; appendicula pyriformi, latiuscula. Mandibula breves, ungue elongato, angusto, falciformi, subtilissime denticulato, apice obtusulo.

Long. 2, lat. 1 millim.

Hab. GUATEMALA, Retalhuleu (*Stoll*).

Body triangular, its anterior margin protracted into a sort of triangular clypeus; colour uniform, scarlet. Body and legs thickly covered with a short velvety pile, out of which spring long, disseminate, pinnate, and slightly curved hairs, which on the shoulders and towards the end of the body are longer and more thickly set than on the remaining parts of the dorsum. The legs are comparatively longer than in *T. mexicanum*, and without brushes on the lower surface. The front tarsus is fusiform, its claws considerably smaller than in the others; the latter are truncate at their top, and bear pinnate hairs, each of which is inserted on an elevation of the integument. The palpi bear elongate, pinnate hairs, which on the second and third joints are arranged in a row; the appendicula is broader than in *T. mexicanum* and bears much shorter hairs. Mandibles short, with a rather cylindrical body, and a long, narrow, slightly curved claw; the concave edge of the latter is, almost imperceptibly, denticulated.

The description and figures are made from living specimens, which were found running amongst dead leaves in the cacao-plantations near Retalhuleu.

3. *Trombidium nasutum*, sp. n. (Tab. III. figg. 1-1 g.)

Corpus fere rhomboidem, depressulum, postice rotundatum, latitudinem maximum inter coxas secundi et tertii pedum paris exhibens, coccineum, tomentoso-hirsutum; tomentum e pilis brevibus quadrifidis constans; pars cephalothoracis frontalis in apicem carnosum producta, clypei instar basem palporum obtegens; coxae lateribus corporis infixae; utriusque lateris coxae anteriores inter se valde approximatae, ab coxis posterioribus (inter se approximatis) ejusdem lateris valde distantes; dorsum impressiones plures transversuales exhibens. Primum et quartum pedum par secundo et tertio paululum longiora; tarsus primi paris pedum fusiformis, unguibus duobus minimis armatus. Palpi marginem clypei superantes; appendicula lanceolata, setas longiores pinnatas gerente. Mandibulae breves, falcis brevis, lata, intus subtilissime denticulata armata.

Long. 1, lat. 0.5 millim.

Hab. GUATEMALA, Retalhuleu (*Stoll*).

Body rhomboidal, broadest between the coxae of the second and third pairs of legs; scarlet, velvety; the frontal part of the cephalothorax forming a sort of fleshy clypeus, which covers the basis of the palpi; a few transverse furrows run across the dorsal surface of the living animal. The coxae are inserted on the sides of the body; the first and second pairs, which are grouped near together, widely separated from the posterior coxae formed by the third and fourth pairs of legs. The first and fourth (!) pairs of legs are longer than the second (!) and third, each thickly covered with short velvety hairs. Front tarsi spindle-shaped, bearing two very small claws on the top; the other tarsi obliquely truncate, their claws almost simple. The palpi bear long pinnate hairs, which on the third and fourth joints are arranged into rows; the appendicula is lancet-shaped and bears on its surface several long setaceous hairs, which, however, are rather shorter than in *T. mexicanum*. The mandibles are short and terminated by a short, broad, falciform claw, the inner edge of which is denticulate.

This species is found amongst dead leaves in the forests round Retalhuleu. The description and figures were made from the living animal.

4. *Trombidium quinque-maculatum*, sp. n. (Tab. IV. figg. 1-1 c.)

Corpus oblongo-ovoidum, depressum, tomentum; nigrum, maculae quinque albis dorsalibus: harum anteriores pares, magnae, triangulares, regionem humeralem utriusque corporis lateris occupantes; reliquae tres maculae impares, scilicet formantes longitudinalem in dorso medio; ex his anterior parvula, rotunda, in dorsi centro sita; huius proxima macula impar magna, ovalis, in abdomine medio sita; in abdominis apice macula magna transversa impar. Apex frontalis cephalothoracis lineolam longitudinalem, brevem, sericeam insuper exhibens. Pedum par primum et quartum longa, crassiuscula, secundo et tertio pari multo longiora, flavescentia, ab articulo tertio usque ad sextum nigrescentia. Palpi flavescentes, pilis nigris sparsis induti; appendicula lata, late inserta. Mandibulae?

Long. 2, lat. 0.75 millim.

Hab. GUATEMALA, near the city (*Stoll*).

Body oblong, a little depressed, with a longitudinal furrow on each side; velvety from closely-set, short, thick, quadrifid hairs: colour deep black, with five white spots; these spots are arranged as follows—two, lateral, large and triangular, occupying the region between the bases of the second and third coxae; a small round one nearly in the centre of the dorsal surface between the apices of these; a larger ovoid one, placed between the centre and the end of the dorsal surface; and a large transverse one on the posterior end. The first and fourth pairs of legs are very long, about the length of the body, considerably longer and stouter than the second and third pairs; their colour is ochraceous-yellow, which from joints 3-6 merges into blackish, owing to the short, pinnate, black hairs by which this part is clothed. Palpi ochraceous.

I found one specimen of this richly-coloured species amongst shrubs in a ravine near the city of Guatemala; it was running about in the bright sunshine. Unfortunately, I accidentally lost the mounted preparation of the mandibles before I had drawn and described them.

5. *Trombidium guayavicola*, sp. n. (Tab. II. figg. 2-2 c.)

Corpus oblongum, tomentosum: nigrum, maculis duabus albis; anteriori in medio dorso sita semilunari, impari, posteriori anali triangulari. Oculi rubri. Pedes ochraceo-rufescentes; per primum et quartum secundo atque tertio longiora, crassiora; tarsorum ungues valde recurvi, ad basin serrati. Palporum articuli cylindrici; appendicula brevis, recta, apice rotundata. Mandibulae?

Long. 1, pedibus extensis 4, lat. 0.5 millim.

Hab. GUATEMALA, Retalhulén (Stoll).

Body oblong; velvety-black, with two large white spots—a semilunar one on the dorsum between the coxae of the third pair of legs, and a triangular one at the posterior margin of the body. Eyes red; the palpi and legs of a clear reddish-brown colour. The first and fourth pairs of legs longer and thicker than the second and third pairs; front tarsi spindle-shaped; claws of the tarsi strongly curved, with some indentations at the basis of their concave edge. Palpi straight, slender, with almost cylindrical joints; the fourth joint with a short claw, and a short, straight appendicula, the latter rounded, very broadly inserted, and bearing several long hairs on the top. The fourth pair of tarsi bear a sort of brush beneath, formed of long, oblique, slightly pinnate hairs: these hairs do not fall off so easily as from the other tarsi, and undoubtedly help the animal in running up the trees.

This species is not rare in the forests near Retalhulén, where it is to be found running up and down the smooth trunks of the guayava trees (palo volador).

6. *Trombidium trilineatum*, sp. n. (Tab. I. figg. 2-2 c.)

Corpus oblongum, pedice rotundato-truncatum, depressiusculum, coccineum, tomentosum, hirsutie brevi, canescenti; pseudo-prothorax albescens, sulco profundo transverso post oculos ab abdomine separatus; dorsum abdominalis hirsutiae albescens, per sulcos duo longitudinales, laterales, parallelos, serpentes in partes tres distinctas separatum. Pedes, palpi et mandibulae rufescentes. Palpi elongati; tertio articulo cylindrico; appendicula brevi, basi lata inserta, apice rotundata.

Long. 1.25, lat. 0.75 millim.

Hab. GUATEMALA, Antigua (Stoll).

Body oblong, with almost parallel sides; the dorsal surface divided into a pseudo-prothorax and an abdomen by a deep transverse furrow, from which two lateral longitudinal furrows take their origin; these latter divide the back of the abdomen into three separate, whitish areas. Legs long, slender, reddish-brown. Palpi slender, clothed with long, stiff, dispersed hairs; their third joint long, almost cylindrical; the appendicula short, rounded at its end, and with a few stiff hairs on the top.

This species lives in the hedges and gardens of the valleys of Antigua and Guatemala city. It is commonly found on the leaves of bushes, where it seems to feed on Aphides. When the specimens, by rubbing themselves against the leaves, &c., begin to lose their whitish velvety pile, their colour appears much redder than in the specimen figured.

7. *Trombidium albicolle*, sp. n. (Tab. I. figg. 3, 3a.)

Corpus oblongum, humeris protractis, apice anali rotundato; coxae humerale, maculis atque striis albis indutum; in apice frontali pseudo-prothoracis macula alba; stria transversa lata inter humeros, postice linguam latam medianam in abdominis dorsum emittens; apex analis albus; duo maculae albae latero-marginales ante corporis apicem sitae, parvae. Coxae lateribus corporis infixae. Pedes longi, graciles, rufescentes, inter se fere aequales. Palpi graciles, pilis longis rarioribus instructi; tertio articulo elongato, cylindrico; quarti unguis unidentato, appendicula recta, apice rotundata, ad apicem piligena.

Long. 0.75-1 millim.

Hab. GUATEMALA, Antigua (*Stoll*).

Body oblong, rounded behind, the shoulders protruding; pseudo-prothorax triangular: colour scarlet, with white spots formed by white, thickly-set, velvety hairs; these spots are arranged as follows—a round white one on the frontal surface of the pseudo-prothorax, a large T-shaped one on the middle of the dorsal surface behind the eyes, an anal one at the end of the body, and two small round lateral ones near the margins of the posterior part of the abdomen. Legs, palpi, and mandibles of a clear reddish colour. Legs long, slender, the first and fourth pairs a trifle longer than the second and third. Third joint of the palpi long, cylindrical; the fourth joint with a small tooth on the concave side of its claw, the appendicula straight, parallel-sided, rounded at the end, where some stiff hairs are inserted.

This species is found with the preceding, on bushes in hedges and gardens in the valley of Antigua.

8. *Trombidium muricola*, sp. n. (Tab. II. figg. 3-3b.)

Corpus breve, obovatum, convexum, humeris rotundatis, apice frontali rotundato; cuticula laevi, sericeo-nitida, nigra, maculis et striis albis, variantibus figuris. Pedes, palpi et mandibulae rufescentes. Pedes longi, graciles; primum par ceteris longius.

Hab. GUATEMALA, Antigua (*Stoll*).

Body short, ovoid, convex, with the shoulders and the frontal and anal apex rounded; smooth, silky-black, with white spots and stripes varying in form in the different specimens: in some the dorsal surface of the abdomen bears a white triangular spot on its posterior third, and from the acute angle of this, which is directed forwards, two narrow stripes proceed obliquely towards the humeri; in others there only remains a small white spot at the anal end, and a narrow white streak running across the dorsum at a short distance behind the middle, the streak in its centre forming a large angle (sometimes connected with the anal spot by a narrow white line) which opens towards the frontal apex; finally, there occur specimens in which the whole body is black, except a small white border along the posterior margin. The legs are long and slender, the front pair a little longer than the others. Legs, palpi, and mandibles light reddish-brown.

This species is found in the rainy season on the adobe-walls of the nopal-gardens

(small plantations of *Opuntia*, upon which the cochineal insects are reared) round the city of Antigua. It runs busily about in the open sunshine.

RHYNCHOLOPHUS.

Rhyncholophus, Dugès, Rech. s. l'ordre d. Acar. en gén. et la fam. des Tromb. en part., Prem. Mémoire, in Ann. des Sciences nat., Zool. i. p. 15 (1834).

1. *Rhyncholophus erinaceus*, sp. n. (Tab. IV. figg. 2-2 d.)

Corpus oblongum, convexum; humeri rotundatis, paululum prominentibus; cuticula molli, non refulgente, ex fusco cinerea, pilis clavatis brevibus, quadrifidis, nigris haud sparse induta; subtus macula alba prope anum. Palpi, mandibulae atque pedes rufescentes, piligeri. Palpi articulo extremo longo, curvo; appendicula longa, lanceolata atque apice rotundata. Epimera primi et secundi pedum paria utriusque lateris in medio corpore valde approximata, per labium fere contigua, ab epimeris posteriorum parium valde distantia; mango corporis anterior desuper insipienti globulum piligerum præbens.

Long. 3, lat. 1-5 millim.

Hab. GUATEMALA, Antigua (Stoll).

Body oblong, not depressed, with somewhat prominent shoulders, the skin not shining, greyish-brown; with short, black, quadrifid, thick and slightly curved hairs, which are not very thickly set, so that the skin can be easily seen between them; the under surface bearing a whitish spot in the anal region. The palpi, mandibles, and legs are reddish-brown, and bear short, black, appressed, obtuse, quadrifid hairs on their upper surface, and longer, imperceptibly pinnate, acute setæ beneath. The fourth joint of the palpi forms a curved tooth, and bears long setæ like the preceding joints and the appendicula; the latter is obtusely lanceolate in form. When looked at from above, the anterior margin of the body appears to be prolonged into a reddish globe, the latter bearing some long black hairs. The apex of the labium forms a sort of flat cup with many marginal fringes.

Two specimens, both of which were found under stones in the neighbourhood of Antigua.

Fam. ACTINEDIDÆ.

Acaride tracheatæ corpore subtriangulari angulis rotundatis, brevi, minime longiore quam latiore, integro. Oculi duo laterales, a margine corporis anteriore valde distantes, facillime perspicui. Cuticula molli, setis rarioribus acutis, subtilissimis pinnatis, induta. Pedes laterales, articuloorum seniorum, epimeria pedum in quoque latere valde approximata. Pedes inter se fere æquales longitudine, setis raris erectis et hirsutis brevi, spissa, appressa induti. Articulus pedum tarsalis gracilissimus, acute terminatus, unguibus duobus curvis armatus; ad eorum basin ungula tertius spinis setulosus insertus est. Palpi quaternorum articuloorum; articulus basalis brevis, secundus longus, quam ceteri crassius, subcylindricus, tertius brevissimus, ad apicem interne dentibus trinis, rectis, obtusis armatus, ultimus quam secundus brevior atque angustior, apice rotundatus, setiger. Mandibulae longæ, ex basi lata sensim apicem versus gracilescentes, in apice unguicula falciformi armata. Inter mandibularum bases in utroque latere corporis nascitur organum longum tubuliforme, angustum, in apice clavatum, quod horizontaliter palpi basin transgrediens marginem frontalem juxta palporum basin ita superat, ut a desuper insipienti distincte et facillime discernatur. Epistoma triangulare, acutum, apice bifido.

ACTINEDA.

Actineda, Koch, Uebersicht d. Arachn. Syst. 3^{re} Heft, 2^{re} Abth. p. 57 (1837).

1. *Actineda flaveola*, sp. n. (Tab. V. figg. 1-1 δ.)

Corpus rotundo-ovoideum, supra convexulum, haud longius quam latius; colore citreo, intestinis albidis translucetibus.

Long. 0.5 millim.

Hab. GUATEMALA, Antigua (*Stoll*).

Body globose, the dorsal surface convex; vivid yellow in colour, with some indistinct whitish spots in the middle of the dorsum, caused by the intestinal contents shining through the semi-transparent skin.

This species occurs on the hills round Antigua, amongst the grass.

2. *Actineda antiguensis*, sp. n. (Tab. V. figg. 2-2 c.)

Corpus breve, subtriangulare vel subpentagonum, paululum depressum, minime longius quam latius; coccineum, maculis albidis indistinctis ex intestinis translucetibus; in dorso sulci tres transversi breves inaequidistantes perspicui; corporis margines laterales antice convergentes, ante oculos angulatum flectentes ad marginem frontalem formandum; marginis posterior late truncatus, angulis rotundatis; hirsuties alba corporis et pedum ita ac in specie precedenti disposita. Tarsorum unguicula stylo elongato inserta, curva, integra, ad eorum basin utrinque seta pinnata oblique inserta; unguis tertius apicem in carunculum imbutiformem dilatatus.

Long. 0.75 millim.

Hab. GUATEMALA, near the city, Antigua (*Stoll*).

Body short, obtusely pentagonal, slightly depressed from above, a little longer than broad; carmine-red, with a few indistinct whitish spots on the dorsum; on the latter are three short, transverse furrows, the anterior of which is situated somewhat behind the eyes, the two hinder ones very near each other in the posterior third of the abdomen; the anterior part from the eyes forward triangular; the front margin rounded; the side margins not parallel, slowly diverging towards the hind one, which is broadly truncate, with rounded angles; on the dorsum are several rows of impinnate white shining hairs. Legs covered with short, thinly set, appressed hairs, amongst which many long, stiff, squarrose setae are conspicuous; the claws are inserted on a sort of petiole, and bear on each side at their base a fringed short seta; false claw forming a cup-like caruncle.

This species is found on shrubs in the vicinity of Antigua and the city of Guatemala; it has a habit of running rapidly up and down the branches of small trees.

3. *Actineda retalteca*, sp. n. (Tab. V. figg. 3-3 c.)

Corpus breve, subtrigonum, antice rotundatum, postice truncatum, angulis lateribus obtusis, paululum depressum; colore rufa, maculis dorsualibus brunneis ab intestinis translucetibus; in medio dorso inter oculos latero-marginales macula trigona; post oculum utriusque lateris stris brunneis longis, antice bifurcatis, oblique ad marginem aulem tendens. Tarsorum petiolus unguiferus nodulis tribus dorsualibus.

Long. 0.75-1 millim.

Hab. GUATEMALA, Retalhuleu (*Stoll*).

Body, legs, palpi, and epistoma reddish-yellow. Body short, indistinctly triangular; anterior margin rounded, the posterior one truncate but with rounded angles; on the middle of the dorsum between the eyes is a triangular brown spot, and behind each eye another long and narrow one which goes obliquely to the poster or margin; these three spots are caused by the intestinal contents shining through the soft and semi-transparent skin. The petiole which bears the claws has three nodules on its back.

This species lives in the woods of the low country about Retalhuleu.

Fam. TETRANYCHIDÆ, Kramer.

TETRANYCHUS.

Tetranychus, Dufour, *Annales des Sciences nat.* xxv. p. 276 (1832).

1. *Tetranychus guatemalæ-novæ*, sp. n. (Tab. VI. figg. 1-1 c.)

Corpus oblongum, antice late rotundatum, postice sensim attenuatum; humeris rotundatis, haud protractis; oculi latero-anteriores; cuticula mollis, semipellucida, subtilissime dense plicatula. Palpi conoidei, crassi, trinoxum articularum. Mandibularum in unam conseruatarum unguicula in setas longas antice convergentes transformatae sunt, basi recurvæ. Tarsi unguibus duobus, inter ambulacra quatuor sitis.

Long. 0.76 millim.

Hab. GUATEMALA, near the city (*Stoll*).

Body oblong, its anterior margin broadly rounded; shoulders not or very little protruding; skin semi-transparent, whitish, very finely wrinkled, with some long, regularly disposed setæ amongst many short ones. Palpi short, rather thick, three-jointed, the last joint with a sort of short double claw. The claws of the mandibles transformed into two thin setæ which converge anteriorly. Labium short, bifurcate. Tarsi with two claws amongst four setæ ('ambulacra' of authors), which bear on the top a small, globose bulb.

This species lives in the vicinity of the city of Guatemala, on a common shrub of the genus *Cassia*; it covers the lower surface of the leaves with its silky webs; the yellowish, comparatively large eggs are protected under circular, transparent covers.

N.B.—I regret that I am unable to offer the reader a more exact and complete description of this species, but I discovered it when preparing to leave Guatemala. I had no time left then for the further study of this interesting species, and I would not have reproduced here the above hasty notes and figures, were it not to prove the existence of the genus *Tetranychus* in Guatemala.

Fam. HYDRACHNIDÆ.

[C. J. Neuman, Kongl. Sv. Vet.-Ak. Handl. xvii. no. 3, pp. 16 et seqq. (1880) (Hydrachnides).]

ATAX.

Atax (J. C. Fabricius), Neuman, Kongl. Sv. Vet.-Ak. Handl. xvii. no. 3, p. 20.

1. *Atax alticola*, sp. n. (Tab. VII. figg. 1-1 g.)

2. Corpus obovatum, satis altum, vix depressulum, antice et postice retundatum, postice tuberculis duobus setigeris prominentibus, albo-flavescent, cuticula transparente; macula dorsalis magna, nigra, margine lobato, valde distincta, per glandulam dorsalem bicusatam in maculas quinque desupra inspiciente divisa; glandula dorsalis antice brunnea, albo marginata, postice flavescent, striam latam medianam longitudinalem formans, ex qua antice et postice rami hinc laterales nascuntur, ita ut glandula bicusata appareat; rami antici latiores, postici angustiores, ramuli irregulares lobiformes emittentes. Pedes et palpi longi, pedum par primum ceteris paulo crassius; palporum articulus extremus tridentatus. Lamine genitales subovales, late; stigmatibus circiter tricornis instructis.

Long. 1 millim.; lat. 0.75 millim.

Max latet.

Hab. GUATEMALA, near the city (*Stoll*).

Body ovoid, very little depressed, transparent, whitish-yellow; the middle of the back occupied by a large black patch which is divided by the dorsal gland into five distinct spots; the dorsal gland forming a broad longitudinal stripe, brown with whitish margins in its anterior half, yellowish behind; this stripe emitting two lateral branches from its anterior third and two also from its posterior third, the anterior of which are comparatively broad and offer several ramifications, whilst the posterior branches are narrow and bear but a few ramifications. Legs and palpi long, slender, transparent, light greenish. The front legs, which are only a little thicker than the rest, bear a few pairs of long stiff spines (these spines being obliquely serrate towards the apex), and their tarsal joint is furnished with a row of short, acute spines; the second pair has only a few short hairs on the dorsal surface, the lower surface showing several pairs of long spines like those of the first pair; the third pair has on the lower surface numerous pairs of squarrose spines and a short pinnate bristle at the apical end of the fifth joint, whilst the apex of the fourth joint bears a tuft of long swimming-hairs; the fourth pair shows on the lower surface of the fourth, fifth, and sixth joints a row of short broad spines, like the teeth of a comb, and the apices of the joints bear, except in the sixth, a more prominent pinnate bristle and tufts of long swimming-hairs. The palpi bear a few stiff spines; and the obtuse top of their fifth joint is tridentate.

This species lives in ponds in the vicinity of the city of Guatemala.

2. *Atax septem-maculatus*, sp. n. (Tab. VIII. figg. 1-1 e.)

Corpus ovatum, convexum, pellucidum exceptis maculis dorsalibus, cum pedibus et palpis ex brunneo albescens; macula dorsalis nigro-fusca, magna, per glandulam dorsalem in maculas septem dijuncta;

BOL. CENTR.-AMÉR., Arachn. Acar., June 1887.

c*

glandula dorsalis lutea, antice lata, quadrifurcata, postice ramulos duos angustos laterales emittens. Pedum par primum ceteris hand crassius, per ultimum ceteris longius. Lamine genitales parvæ, se invicem angulis internis tangentes, obtuse pentagonæ; stigmatibus majoribus quinis, minoribus bis instructæ.

Long. 0.5 millim.

Hab. GUATEMALA, near the city (*Stoll*).

Body ovoid, convex, transparent; brownish-white, like the legs and palpi; the dorsal patch large, blackish-brown, separated into seven single spots by the orange-coloured dorsal gland; the latter divided in its anterior half into two bifurcate branches, whilst at its posterior end it emits two small and narrow lateral branches. The first and second pairs of legs bear on their lower surface a few pairs of long, squarrose spines, which are indistinctly serrate at the end and inserted behind tooth-like elevations of the epidermis; the third pair with a pinnate bristle at the apical end of the fifth joint, and also with several swimming-hairs beneath; the fourth pair with a few short spines beneath, the spines gradually becoming longer towards the apex of the joints, and the apical end of the third, fourth, and fifth joints with a short, pinnate bristle, and towards the apices of the same joints some tufts of long swimming-hairs. The genital plates are small, situated in the middle between the epimera of the fourth pair of legs and the abdominal margin, and touch each other at their inner angles; each plate bearing five large and two small stigmata.

Found in ponds in the vicinity of the city of Guatemala.

Var. ypsilon. (Tab. IX. figg. 1-1 c.)

Corpus oratum, pellucidum; macula dorsali magna nigro-fusca, granulosa; glandula dorsalis lutea, Y-formis, integra. Lamine genitales pentagonæ, parvæ, angulis internis se invicem tangentes, stigmatibus quinis vel senis instructæ. Pedes atque palpi sicut in typo.

Long. 0.25 millim.

Hab. GUATEMALA, near the city (*Stoll*).

Body ovoid, transparent, with a large brown dorsal patch, which is divided by an orange-coloured Y-shaped dorsal gland into three distinct spots. Genital plates pentagonal, touching each other at their anterior angles, each bearing five or six large and two small stigmata. Legs and palpi as in the type.

Found in the same pond with the type in the vicinity of the city of Guatemala.

I cannot consider this little *Atax* specifically distinct from *A. septem-maculatus*, with which it agrees in the colouring, in the shape of the legs and palpi, and in the character and size of the genital laminae. It differs from typical *A. septem-maculatus* in being a trifle smaller, and in the form of the sharply marked, non-ramified dorsal gland.

3. *Atax dentipalpis*, sp. n. (Tab. X. figg. 1-1 d.)

Corpus oratum, fere rotundum, depressulum, pellucidum, flavo-albescent, postice tuberculis duobus setigeris; macula dorsalis nigra, lata, per glandulam dorsalem brunneam, albo-marginatam, longam, antice bifur-

catam divisa. Pedes longissimi, setigeri, secundo et quarto pari ceteris longioribus. Palpi elongati, apice tridentato, obtuso; articulo quarto longo, cylindrico, cum dente longo valido, crasso in medio, armato. Lamine genitales disjunctae, latae, arcuatae; stigmatibus senis, in binos aevros trinostrum stigmatum segregatis instructae.

Long. corp. 0.75 millim.; long. pedum 1.5 millim.

Hab. GUATEMALA, near the city (*Stoll*).

Body ovoid, almost globoid, a little depressed, transparent, brownish-white, with two setigerous tubercles at its posterior end and (in the male) two small tufts of hairs at the end of the genital fissure; dorsal patch black; dorsal gland Y-shaped, brown, with white margins, dividing the dorsal patch into an anterior triangular, and two long, broad, lateral spots. Legs very long and slender, the second and fourth pairs longer than the first and third, all bearing numerous long hairs and rows of short spines. Palpi very long; fourth joint cylindrical, bearing on the middle of its inner surface a strong, obliquely inserted tooth which points forward; fifth joint long, slightly arcuate, with an obtuse tridentate apex. Genital plates situated near the abdominal margin, arcuate; each bearing six large stigmata, which are divided into two equal groups. The epimera of the third and fourth pairs of legs form an almost quadrate plate.

Lives in stagnant ponds in the vicinity of the city of Guatemala.

NESSEA *

Nessea (C. L. Koch), Neuman, Kongl. Sv. Vet.-Ak. Handl. xvii. no. 3, p. 29.

1. *Nessea guatemalensis*, sp. n. (Tab. X. figg. 2-2 ♂, ♀; and Tab. XI. figg. 1-1 f, s.)

♀. Corpus ovatum, convexum, pellucidum, albido-flavescent; macula dorsali fusco-nigrescente, per glandulam dorsalem angustam, antice bipartitam, luteolam in quinque maculas disjunctas; tres anteriores minores, breves, duas posteriores majores, saepeque exterior arcuatas. Pedes medietate longitudinis, a primo pari gradatim longitudine crescentes, unguiculis valde recurvis; pedes articuli quarti etque quinti fasciis setarum in seriem obliquam dispositarum ad apicem instructi; pedes posteriores spinarum serie brevium subrectis armati. Palporum articulus secundus crassus, quartus dente armatus. Oculi duo magni, conici, divergentes. Area genitales stigmatibus numerosis, circa 18 magnis etque 8 parvis instructa, quae aream semilunarem in utroque latere fissurae genitalis occupant.

Long. 0.75 millim.

♂. Corpus ovatum, altum, antice obsolete truncatum; macula dorsali fusco-nigrescente in dorsi dimidio posteriore sita, brevior quam in femina. Mandibula versus apicem serrulata. Pedum unguiculus valde recurvus, basi incrassato inserto; pedum quatuordecim articulus quartus in medio excavatus, seriebus duabus dentium oblique oppositis in marginibus excavationis armatus. Area genitales stigmata diffuse disposita, circiter viginti.

Long. 0.5 millim.

Hab. GUATEMALA, near the city (*Stoll*).

2. Body ovoid, convex, whitish-yellow, transparent; dorsal patch brownish-black; dorsal gland narrow, yellowish, anteriorly bifurcate, and dividing the dorsal patch into

* This genus requires a new name; it is preoccupied in Polypi (Lamouroux, 1812) and Mollusca (Risso, 1826).

five distinct spots—three short ones before, and two large, longitudinal, parallel ones behind the lateral branches. Legs of moderate size, the first pair the shortest, the following pairs gradually longer; all bearing tufts of swimming-hairs and spines of varied length, and the fourth and fifth joints at their apical ends with tufts of long swimming-hairs arranged in oblique rows; the fourth pair with a row of short spines on its lower surface; the claws strongly curved, forming an acute angle. Palpi with the second joint thicker than the rest, the fourth joint bearing on its middle an oblique setigerous tooth. Eyes conical, diverging. Stigmata of the genital area numerous, about twenty-six on each side, arranged on a semilunar space.

♂. Body somewhat truncate at its anterior margin; dorsal patch limited to the posterior half of the back, the anterior half milk-white. Claws strongly curved, both of them inserted on a common swollen basis. Stigmata of the genital area numerous, about twenty in number, not arranged in a semilunar area as in the female, but rather irregularly disseminate on an obliquely transverse, oblong space. The fourth joint of the hind legs bears on the middle of its inner side an excavation, the margins of which are beset with rows of obliquely set spines or teeth placed opposite to each other. The concave edge of the mandibles shows some very small indentations towards the apex.

The male and female above described were found in a pond in the vicinity of the city of Guatemala. Not having seen them *in copula*, nor bred them, I am not quite sure that they belong to one and the same species; yet from their general appearance, their common habitat, and from the fact that I found no other *Nesaea*, except *N. numulus*, in this pond, I am inclined to think that the above-described specimens really belong to the same species.

2. *Nesaea numulus*, sp. n. (Tab. XI. figg. 2-2 c.)

2. Corpus orbiculatum, nummuli instar valde depressum, antice truncatum, in margine anteriore utrinque tuberculis trias piligeris juxta oculum instructum; colore fuscescente, opaco, cuticula cribelli laeviter perforatula; macula dorsalis lata, margine posteriore lobata, nigra, in dimidio anteriore maculis clavis pennis rotundis interrupta, in medio per glandulam dorsalem, latam, fuscam, postice utrinque dilatata divisa. Pedes breves, a primo pari gradum longitudine crescentes, setigeri, piligeri; per quantum serio spinarum atque seta pinnata ad articuli quarti et quinti apicem subtilis instructum; articulus extremus ad apicem subtilis subexcavatum. Palpi marginem corporis anteriorem paululum superantes, tenues; articulo quarto arcuato, dentigero. Stigmata areae genitalis numerosa, circiter triginta in quoque latere. Oculi in margine anteriore siti.

Long. et lat. 0.75 millim.

Hab. GUATEMALA, near the city (Stoll).

Body orbicular, very much depressed, truncate at its anterior margin, the latter bearing three small tubercles on each side (under the microscope the skin seems to be perforated by densely set little holes, like a sieve); colour brown; dorsal patch black, broad, with several transparent brown spots on its anterior half, its exterior margins lobate; dorsal gland compact, broad, brown, truncate before, laterally extended behind. Legs short, the first pair the shortest, the following pairs gradually longer; all

bearing a number of spines of varied length and numerous swimming-hairs, the latter being longest in the fourth pair; hind pair with the fourth and fifth joints bearing a row of erect short spines (the most apical of which are pinnate) beneath, and the lower surface of the tarsal joint slightly excavated at the apex. Stigmata numerous, about thirty on each side, disseminate on both sides of the genital plate.

This species lives with the preceding in the vicinity of the city of Guatemala.

LIMNESIA.

Limnesia (C. L. Koch), Neuman, Kongl. Sv. Vet.-Ak. Handl. xvii. no. 3, p. 97.

1. *Limnesia guatemalteca*, sp. n. (Tab. VII. figg. 2-2 e.)

Corpus ovatum, altum, pellucidum, late fuscescens, obscure punctulatum maculis dorsualibus quinque fuscis: tribus anterioribus, duobus (aliquando tribus) posterioribus, glandule dorsualis albescentis ramulis divisis. Oculi valde distantes, bini in quoque latere inter se vicini. Palpi et pedes fuscescentes. Pedum par primum ceteris crassius, paria trina anteriora articulis extremis oblique truncatis, unguiculatis; quarto par angustius carente, omnia sparse setigera, spinigera. Lamine genitales parva, ab epimeris posticis trigonice distantes; margine inferiore recto se invicem tangentes, exteriori sinuato; binis stigmatibus in utraque lamina longitudinaliter dispositis. Mandibule: ungues falciformi integro. Palporum articulus extremus apice tridentatus.

Long. 0.5 millim.

Hab. GUATEMALA, near the city (*Stoll*).

Body ovoid, high, transparent, light brown, with a fine dark punctulation; dorsal patch divided into five brown spots—three before, two behind—amongst which the whitish dorsal gland spreads its ramifications, the limits of the latter not being so distinct as in many other species. First pair of legs shorter and a little thicker than the others; the tarsal joint of the three anterior pairs obliquely truncate at the apical end. Genital plates small, placed at a little distance behind the posterior epimera, and touching each other along the entire length of their interior margin, their side margins slightly sinuate and diverging; each lamina bearing two large stigmata only. Mandibles with a falciform, narrow, non-denticulated claw.

This species lives in ponds near the city of Guatemala.

2. *Limnesia longipalpis*, sp. n. (Tab. IX. figg. 2-2 c.)

Corpus ovatum, pellucidum, altum; macula dorsuali nigra, triloba; glandula dorsuali sulphurea—antice bifurcata, ramis lateralibus valde distinctis, angustis, lobulatis; postice lata, minus distincta. Pedes longitudinalibus mediocribus, a primo pari gradatim longitudine crescentes; tris paria anteriora sparse setigera, articulis tarsalibus oblique truncatis; ultimum par subitus serie setarum natatricium longum instructum. Palpi valde elongati; secundo articulo brevi, crasso, dente brevi interne instructo; quarto articulo longissimo, dentigero; ultimo armato, obtuso, tridentato. Lamine genitales angustae, ab epimeris posticis distantes; margine interno se invicem tangentes; marginibus externis divergentibus; trinis stigmatibus, uno antice solo, duobus postice vicinis, instructae.

Long. 0.5 millim.

Hab. GUATEMALA, near the city (*Stoll*).

Body ovoid, transparent, high, whitish-yellow, with a black, trilobate dorsal patch; dorsal gland sulphur-yellow—anteriorly bifurcate, and with narrow branches and sharply marked outlines; posteriorly broad, obfuscated, and with rather indistinct outlines. Legs of moderate size, gradually longer from the first to the fourth pairs; tarsal joint of the first three pairs bearing several long hairs and spines, and obliquely truncate; the fourth and fifth joints of the hind legs with a row of long swimming-bristles beneath. Palpi exceedingly long; the second joint rather stout, and with an erect, short tooth; the fourth joint very long, slightly curved; the fifth joint with an obtuse, tridentate top. Internal edge of the falx of the mandibles very finely denticulated. Genital plates narrow, placed at a little distance behind the hinder epimera, and touching each other along the entire length of their interior margin; their exterior margins diverging; three large stigmata on each lamina.

Lives in ponds in the vicinity of the capital of Guatemala.

3. *Limnesia puteorum*, sp. n. (Tab. VII. figg. 3-3 c.)

Corpus ovatum, altum, opacum, fuscum; maculis dorsualibus nigris, irregularibus; glandula dorsalis compacta—antice alba, longe bifurcata; postice brunnea, indistincta. Pedes medioeris longitudinis, unguiculis ante apicem dente instructis; antici sparse setigeri, postici (paris tertii et quarti) longis setis natatricibus instructi. Palpi longi, articulo secundo unidentato. Mandibulae falci integrae. Laminae genitales oblongae, margine interno recto, inter se valde approximatae; stigmatibus trinis magnis instructae.

Long. 1.0 millim.

Hab. GUATEMALA, Antigua (*Stoll*).

Body ovoid, high, not transparent, brown, with irregular black spots and stripes on its back; dorsal gland white and very distinct, bifurcate in its anterior half, brown and indistinct behind. Legs of moderate size, the hind pair with rows of swimming-hairs; tarsal claws bearing a tooth on their concave edge. Palpi long, the second joint with a tooth on its inner side. Claw of the mandibles not denticulated. Genital plates oblong, narrow, touching each other along the entire length of the interior margin; each bearing three large stigmata.

Lives in the "pillars" (water-cisterns) of the city of Antigua.

4. *Limnesia læta*, sp. n. (Tab. VIII. figg. 2-2 d.)

Corpus ovatum, fere globetum, altum, pellucidum; maculis nigris dorsualibus trinis in utroque latere—binis internis magnis, singula externa parva; medio in dorso stria lata longitudinalis lutescens, antice, postice atque in medio transverse nigrescens. Pedes haud longi, a primo pari gradatim longitudine crescentes, setis atque spinis variis instructi; per quantum subius serie spinarum erectarum brevium atque setis natatricibus longis indutum; in apice articuli tertii et quarti ejusdem paris seta pinata adest brevis. Palpi breves, articulus quartus dente setigero armatus obliquo. Mandibulae unguiculis falciformibus, angustis, integris. Laminae genitales in unam, polygonam, antice fissuram brevem lacunculam præbentem conerectae; trinis stigmatibus instructae, epimeris posticis valde approximatae.

Long. 0.5 millim.

Hab. GUATEMALA, near the city (*Stoll*).

Body ovoid, almost globoid, high, transparent; yellowish-white, with six black spots on the back, these being widely separated by a broad longitudinal yellow stripe which at its frontal and anal end and again in its middle is itself crossed by a transverse blackish stripe, the spots placed thus—two large ones on the anterior half of the dorsum and two others on the posterior half, the latter separated from the former by a white interspace, and on each side at a little distance from the anterior ones is a small one nearer to the margin. Legs of moderate size, rather short, gradually longer from the first to the fourth pair; all bearing many spines and swimming-bristles, which are most numerous on the lower surface of the fourth pair; on the latter beneath there is also a row of short, erect spines on the third, fourth, and fifth joints, together with tufts of long swimming-hairs. Palpi rather short, their fourth joint bearing an oblique tooth. Mandibles with a narrow, falciform, non-denticulated claw. Genital plates united into a single large pentagonal piece, which only in its anterior half bears a lanceolate fissure; three large stigmata present on each side of the genital piece.

Lives in ponds in the vicinity of the city of Guatemala.

Fam. BDELLIDÆ.

[Kramer, Grundzüge zur Systematik der Milben, in Archiv für Naturg. xliii. p. 244 (1877).]

BDELLA.

Bdella, Latreille, Gen. Crust. et Insect. i. p. 153 (1806).

1. *Bdella splendida*, sp. n. (Tab. III. figg. 2-2 c.)

Corpus oblongum, leve; parte antica magna, trigona, in rostrum longum acutum porroctum desinens; humeri rotundati, post eos corpus paullo angustius, usque ad apicem analem gradatim attenuatum; portice late rotundatus; color ex fusco rubro. Rostrum ad basin paululum infistum, ad apicem subtus piligerum, setula dimidio paululum separata, divergentes. Palpique rostrum longiores, setis sparsis brevibus armati; ad apicem quinti articuli valde elongati setae duae inaequales ferentes, internam breviorum, externam longiorem. Extremus palporum articulus secundo brevior, tertius atque quartus brevissimi. Mandibulae elongatae, ad basin latiores, apicem versus gradatim gracilescentes, chela minimis instructa; chelarum dens fixus curvus, acutus; dens mobilis latus, apice truncatus. Pars antica corporis (pseudo-cephalothorax), pedes et palpi soccinati; pars postica (abdomen) ex fusco rubra, stris lata roseae at lineis tribus transversis angustis inaequidistantibus ejusdem coloris induta. Cuticula mollis, subtilissime striatula, in dorso setae sparsae in series longitudinales dispositae gerens.

Long. 1-25 millim.

Hab. GUATEMALA, near the city (*Stoll*).

Body oblong, smooth, gradually narrower from the rounded shoulders towards the anal end, the latter broadly rounded; pseudo-cephalothorax triangular, ending in a long, acute rostrum. The pseudo-cephalothorax, rostrum, palpi, and legs are vivid scarlet; the rest of the dorsal surface from behind the eyes to the anus reddish-brown with darker spots, the latter varying with and depending on the intestinal contents; a

broad longitudinal stripe of a rosy light-red occupies the middle of the dorsum, and three unequally placed transverse lines of the same colour run across the back. Palpi longer than the rostrum; the second joint the longest, the third and fourth joints very short; the fifth joint shorter than the second, obliquely truncate at the apical end, where it bears two long stiff setæ, the inner seta being shorter than the outer one. Mandibles very long, broad at their base, narrowing gradually towards the end; chela very small, their immovable tooth acute, falciform, the movable one broad, truncate at the end. Skin soft, showing under the microscope very fine and densely-set wrinkles. The dorsum bears several stiff setæ arranged in two longitudinal rows. The point of the epistoma has a small brush of short hairs beneath, which latter, by a narrow median interstice, are divided into two. The hairs of the palpi and dorsum are only most finely fringed, those on the legs being quadrifid. Claws of the tarsi broad, curved; the false (third) claw with a short-haired brush.

This pretty *Bdella* lives amongst dead leaves in the hedges and gardens of the city of Guatemala.

Note.—On the 10th of July, 1880, I found the larva of an Acarid adhering to one of the fore legs of a large Elateroid beetle, *Chalcolepidius*, sp., in the woods near Retalhuleu, and some days later a stripped-off skin of the same kind of larva on the bark of a tree.

This larva, which I have figured under the doubtful name of *Bdella*, sp. (Tab. III. figg. 3-3d), is 0.5 millim. long, and reddish-yellow in colour, with an ovoid abdomen, which is attenuated towards the rostrum (thus forming a sort of collum, on which a stout rostrum is inserted); the shoulders a little prominent, rounded, and with a large black eye-like spot on each side near the margin; the dorsal surface of the soft abdomen beset with several transverse rows of short, quadrifid, somewhat clavate setæ, and the skin densely and finely wrinkled. The palpi four-jointed: the basal joint short, the second thick, the third narrow, cylindrical, the fourth bearing a falciform claw and a straight appendicula (the latter resembling that of the true *Trombidia*); and with a few pinnate hairs spread over the surface, more numerous hairs adorning the appendicula. Mandibles long, consisting of a large, broad, basal piece, which is attenuated rather suddenly into a long, narrow branch, the latter bearing on its top an extremely small tooth. Legs long, slender, with two curved claws and a pinnate false claw.

Prof. G. Canestrini and Prof. F. Fanzago give in their excellent treatise "Intorno agli Acari Italiani" (Atti Soc. Pad. v. 1877) the figure (tab. 4. fig. 1) and description (pp. 70 *et seq.*) of an Acarid larva, which bears a strong resemblance to the above-described larva from Retalhuleu, from which it differs, however, in the want of eyes and of a false claw. The learned authors are of opinion that their larva is that of *Rhyncholophus electoralis*, Koch, or of an allied form. The species from Retalhuleu has in common with *Rhyncholophus* only the short quadripinnate hairs of the back and

of the fourth joint of the palpi; it differs from the adult *Rhyncholophi* by the general shape of the body, the insertion of the palpi, the configuration of the mandibles and tarsi, and the finely wrinkled skin.

Fam. EUPODIDÆ.

[Kramer, Ueber Milben, Zeitschr. ges. Naturw. liv. p. 448 (1881).]

SCYPHIUS*.

Scyphius, C. L. Koch, Ueberr. d. Arachniden syst. Heft iii. Abth. 2, p. 62 (1837); Kramer, l. c. p. 449 (1881).

1. *Scyphius maniacus*, sp. n. (Tab. VI. figg. 2-2 d.)

Corpus oblongum, antice acutum, postice rotundatum, supra convexulum, albidum, semipellucidum, laeve, setis raris in pedibus et abdomine indutum; humeri late rotundati, post eos abdomen coarctatum. Pedum primum par ceteris longius, per secundum brevius. Pedes corporis margini laterali inserti. Pseudocephalo-thorax triangularis, ab abdomine per sulcum transversum separatum. Oculi nulli. Palpi longi, eorum articulus extremus penultimo longior, cylindricus, basi coarctatus, apice rotundatus, setis pluribus rectis tacitibus armatus. Mandibulae epistomium superantes, cheliferæ, chelarum articulus fixus curvus, dense acuto armatus, ad basin supra setulam gerens; articulus mobilis falciformis, acutus, angustus, basi solum atque extremo apice articulum fixum tangens. Pedum articuli extremi longi, versus apicem gracilescentes, unguicula curvis, unguicula tertia spuria, recta, setulosa armati.

Long. 0.75 millim.

Hab. GUATEMALA, Retalhulen (*Stoll*).

Body oblong, rather acute towards the frontal end, rounded behind, convex, soft-skinned, semitransparent, whitish, smooth, with a few rows of stiff, minutely pinnate hairs on its back. Abdomen a little attenuated behind the rounded and not very prominent shoulders. Legs inserted on the sides of the body, the first pair longer, the second pair shorter, than the rest; epimera of pairs 1 and 2 separated by an interspace from those of pairs 3 and 4; joints cylindrical, except the sixth, the latter gradually attenuated towards the apical end, and bearing two curved claws and an erect brush-like false claw. Legs and palpi with several squarrose bristles. Palpi long, their last joint rounded at the end, and bearing several long, stiff setæ. Mandibles short, but visible from above; the fixed branch of the chela ending in a short curved tooth, which bears a short seta of about the same length at its base outside; the movable branch strongly curved, falciform, narrow, and touching the tooth of the immovable branch only with its acute point; none of the branches bear any denticulations. Eyes wanting.

This species lives in the forests of the "tierra caliente" near Retalhulcu among the fallen leaves. It runs in a very quick but interrupted erratic manner, and is rather difficult to capture. The softness of its skin renders microscopical examination somewhat difficult.

* Preoccupied in *Picces* (Rimo, 1826).

Fam. IXODIDÆ.

[C. L. Koch, System. Uebers. über die Ordn. der Zecken, in Arch. f. Naturg. x. Jahrg. Bd. I, p. 220 (1844).]

IXODES.

Ixodes, C. L. Koch, loc. cit. p. 231.

1. *Ixodes boarum*, sp. n. (Tab. XIII. figs. 1-1e; and Tab. XIV. fig. 4, 2.)

♂ latet.

♀. Corpus ovatum, depressum, glabrum, in superficie dorsali et ventrali abdominis tam siccis longitudinalibus quam punctis impressis instructum; colore, animalculo vivente, ex ceruleo sericeo (in animalculo alcoholo preservato sordide ochraceo). Lamine stigmatice obtuse triangulares, stigmati excentrico, rimam simplicem oblongam formante. Scutum occipitale latum, parvum, angulis obtusis, nitidum, punctis et forosis raris impressis indutum, colore castaneo uniformi. Oculi nulli. Scutum frontale parvum, glabrum, foris frontalibus cavis, colore castaneo. Rostrum parvum, breve, colore late castaneo. Area mandibularis hamulis senis in utroque latere armata, quarto latissimo, quinto longissimo. Area maxillaris subtus dentibus squamiformibus, rotundatis instructa. Palpi breves, rostrum longitudine haud superantes. Pedes graciles, breves, setulis raris armati.

Long. secundum satietatem varians: 2·5 millim. in jejunis, 5 millim. in satiatie animalculis; lat. 3·5 millim.

Hab. GUATEMALA, Retalhuleu (*Stoll*).

Male unknown. Body of the female ovoid, depressed, hairless. The dorsal surface of the abdomen shows several linear furrows, arranged in two or three concentrical series which radiate from an imaginary centre in the occipital plate. The whole skin of the abdomen perforated by a multitude of extremely small punctiform dimples. Stigmatic plates very simply built, triangular, with rounded angles, the stigma forming a simple excentric fissure. The occipital plate is small, comparatively broad, with obtuse angles, shining, of a uniform chestnut-brown colour, with some punctiform dimples, principally on the sides. No eyes visible. The frontal plate small, chestnut-brown, devoid of the two frontal grooves or fovee common to the females of *Amblyomma*. Rostrum small, short. Mandibles bearing six hooks on each side, only five of which are distinctly visible; the fourth is the broadest, the fifth the longest of all. The maxillary teeth are scaly, rounded. The palpi are rather short; the legs comparatively short and slender. The total length of the body varies according to the state of digestion; from 2·5 millim. in the empty specimens up to 5 millim. in the satiated ones. Breadth 3·5 millim.

I found about sixty individuals of this species, all of them females, adhering to the skin of a *Bos imperator*, near Retalhuleu. The rudimentary state of the genital orifice renders it possible that the specimens are not adult.

AMBLYOMMA.

Amblyomma, C. L. Koch, System. Uebers. über die Ordn. der Zecken, in Arch. f. Naturg. 1. Jahrg. Bd. 1, p. 223 (1844).

1. *Amblyomma mixtum*. (Tab. XII. figg. 1-1 ♀; 2-2 ♂, ♂.)

Amblyomma mixtum, C. L. Koch, loc. cit. p. 227; Uebers. d. Arachniden syst. Heft iv. pp. 74, 75, t. 13. figg. 47 (♂), 48 (♀).

♂. Corpus ovatum, valde depressum, sulcus concavum, colore ex rufo brunneo. Superficies dorsalis punctis numerosis impressis atque lineolis arcuatis irregularibus, interruptis, undulatis, flavo-albescentibus ornata, quae ex dorsi centro, radiorum instar, ad peripheriam corporis petunt. Earum dum maximum angulum humeralem cuiusque lateris capessunt inter quas area triangularis fere unicolor et lineolis albidis fere carnis interest. Forae frontales nullae. In abdominis parte posteriore lineola obscure fusca et intestinis translucentibus nata, albidis se adiacent. Inter sulculos marginis posterioris lineola albidæ et fuscae, breves, radiatim per pares alternantes dispositae, adsunt. Scutum frontale, rostrum et palpi rufo-brunnei. Orificium genitale rimam angustam transversalem formans. Lamina analis in utroque latere marginis posterioris setis duabus longis armata. Pedes rufo-brunnei, setulis paucis serieis armati.

Long. 4.5 millim.; lat. 3.5 millim.

♀. Corpus ovatum, in jejunio valde depressum, in satietate depressio-globosum. Scutum occipitale magnum, triangulare angulis rotundatis, centrum dorsi angulo postico fere attingens, laeve, colore rufo-brunneo, marginibus lateralibus fascia, maculis albidis irregularibus, disjunctis, e margine postico in ramos duo ramificatos discedentibus. Forae punctiformes sento impressae, numerosae, nigrae. Abdomen colore fusco-olivaceo, margine clarius brunneo, haud nitidum, setulis albidis raris ornatum, punctis, sulcis et foris impressis inaequum. In jejunio intestina colore fusco translucent, in satietate abdomen colorem uniformem obscure purpureum seu rubidum exhibet. Lamina stigmatica triangularis, angulis late rotundatis, latoribus arcuatis. Rima stigmatica claviformis. Area genitalis triangularis; orificium genitale transversale angustum, margine anteriore subtilissima denticulato. Area analis rectangulata angulis rotundis, valvulis analibus semilunatis; quaeque valvula in apice anteriore setis duabus, in posteriore setis tribus armata. Scutum frontale forae duabus rotundis ornatum. Palpi, rostrum et pedes in ambobus sexibus aequaliter constructi. Palpi rostrum haud asperantes, compressi, articulo quarto brevi, in apice tertii inserto, colore corporis, ad apicem secundi et ad basin tertii articuli macula fusca interne ornati. Apex rostri albidus-flavus. Basis rostri et anguli posteriores scuti frontalis fusci. Area maxillaris colore saccineo, dentibus conoideis, sat erectis, inter se sat distantibus armata. Area mandibularis in utroque latere hamulis quinque armata: eorum primus longissimus, secundum velut appendiculum brevem insertum ferens; tertius, quartus et quintus in ramo segregato dispositi. Pedes rufo-brunnei, setulis paucis armati. Pedes antei longiores, externo denticulati, ceterorum parium articulus quintus in apice dentibus duabus armatus. Variat femina hujus species in singulis individuis colore plus minusve clariore scutorum occipitalis et frontalis, nec non rostri et pedum. Accedit ad hoc quod macula albidæ scuti occipitalis ramificatae extensione multum variant.

Long. 4.5 millim., lat. 3.5 millim. in jejunio, ad long. 12 millim. et lat. 8 millim. in satietate.

Hab. MEXICO (Köck); GUATEMALA, Retalhuleu, Antigua (Stoll); NICARAGUA, Chontales (Janson); COSTA RICA, Cache (Rogers).

♂. Body oval, much depressed, concave on its lower surface, testaceous, the dorsal surface with numerous punctiform black and testaceous dimples and several irregularly shaped, arcuate, narrow whitish stripes, which radiate from the centre of the dorsum and are sometimes interrupted so as to form a mere series of spots; a triangular area behind the anterior margin of the collar is comparatively free from these spots. On each side-margin and parallel to it there runs a narrow longitudinal stripe, which

begins near the white eyes and ends in about the middle of the lateral margin. In the hinder part of the body the intestines shine through the semipellucid skin in the form of blackish spots and stripes. On the interstices between the furrows of the posterior margin there occur alternating pairs of black and white linear stripes. The arrangement of the white colouring is subject to individual variation. Colour and shape of the frontal plate, of the rostrum, palpi, and legs, similar to that of the female, except the two frontal dimples, which only occur in the female and are wanting in the male*. The genital plate bears a transverse narrow fissure. The anal plate bears two setæ on each side of its posterior margin.

♀. Body oval, in empty specimens much depressed, in the satiated ones globose. Occipital plate triangular, almost reaching the centre of the dorsum, shining, testaceous, with dark brown side margins. From the posterior angle, which is yellowish-white, there proceeds a ramified branch of the same colour towards the anterior margin of each side: these branches vary much in the different specimens; in some they are broad and continuous, in others they are narrow and tend to resolve themselves into several spots. The abdomen is dark brown, opaque, lighter at its margin, irregularly dimpled and furrowed, and bears short, thinly set whitish hairs. In transparent light under the microscope the ramifications of the intestines are visible in the form of blackish arcuate stripes. In the specimens which are filled with blood the abdomen assumes during life a uniform dark purple hue. The stigmatic plate in both sexes is triangular, its fissure claviform; the stigma proper presents itself as an arcuate small hole in a dark chitinated lamina. The genital plate is triangular, with a narrow transverse fissure, the anterior margin of which is finely denticulated. The anal valvula of each side shows two setæ on its anterior and three on its posterior end. The front plate bears two round dimples. The eyes are white. The palpi are compressed, similar in colour to the body; they bear at the top of the second and the base of the third joint a small brown spot, and are beset with several short hairs; the fourth joint, which is very small, is inserted centrally at the top of the third one. The rostrum is of a light transparent brown, yellowish at its extremity. The mandibles bear five hooks arranged on two branches, the first bearing the first and second hooks, the second bearing the third, fourth, and fifth hooks; the second hook is very short and forms a sort of small appendage to the first one. The maxillary teeth are conoid, obliquely erect, somewhat distant from each other, amber-yellow. The legs are light brown, whitish at the apex of the joints; the first pair have their last joint irregularly denticulate at its apex, and the other pairs bear two teeth at the apex of the fifth joint.

This species is the most common of all the Ixodidae of Central America, and generally known by the name of "*garrapata*," which is a corruption of "*agarrapata*" (clapping something with the legs). I have never found the male in a parasitic state,

* They have been drawn by mistake in fig. 2 of Tab. XII.

only free on grass and hushes in the "tierra caliente" and "tierra fria" of Guatemala (Retalhuleu, Guatemala city). The female, which abounds in the woods and savanas on grass and hushes, is occasionally rubbed off by horses, cattle, or dogs, and even by man. It adheres tenaciously to the skin, fixing itself by perforating the cutis with its sucking-apparatus; and remains, when undisturbed, for several days, till filled with blood, and then probably falls off spontaneously by its own weight. If forcibly removed, the sucking-apparatus breaks off and remains in the wound, causing a disagreeable and sometimes painful inflammation for a considerable time, but I never saw any serious consequences result from it. Even in its juvenile state the garrapata is of parasitic habits. The young, which are distinguished by the inhabitants of Guatemala by the name of "mostacilla" (derived from "mostaza," mustard), hang to the grass in clusters of thousands, especially during the dry season; and by their creeping on the bare skin and frequent biting they form one of the greatest plagues to the European traveller, who is sometimes kept awake for hours during the night by them. The males I have heard spoken of as "conchuda." The female has been collected by Mr. Janson in Nicaragua, by Mr. Rogers in Costa Rica, and by myself in many places of the "tierra caliente" and "tierra fria" of Western Guatemala (Retalhuleu, Escuintla, Antigua, Guatemala city).

Remarks. Though I have not seen the types of *A. mixtum*, which Koch describes as from Mexico, I cannot doubt that the above described *Amblyomma* really belongs to that species. As Koch describes and figures both sexes, his must therefore have been a common species; and the above described is the most common of all Ixodidae in Central America, and probably also in Southern Mexico.

2. *Amblyomma foreli*, sp. n. (Tab. XII. figg. 3-3♂; and Tab. XIV. figg. 3, 3a-3d, ♀.)

♂ latet.

♀. Corpus ovato-depressum, abdominis cuticula (in exemplo satiato, in alcoholo preservato) ex griseo fusca ac olivacea, punctis, striolis transversalibus atque maculis numerosis fuscis marmorata. Scutum occipitale oblonge triangulatum, angulis rotunde truncatis, planum, nitidum, punctis nigris magnis raris impressis in superficie dorsuali ornatum, nigrum, in angulo postico maculam colore siccineo ferens, e qua stria lata mediana obscure castanea ad marginem scuti anteriorem petens nascitur. Margo humeralis punctis impressis caret. Scutum frontale nigrum, nitidum, punctis impressis raris ornatum; foreole duae frontales oblongae, postice truncatae, antice angulatum protractae. Arca mandibularis basi nigra, apicem versus flavescens, hamulis quinque, binis in primo, ternis in secundo ramo dispositis armata; arca maxillaris colore succineo. Palpi fusci, nitidi, superficie interna deplanati, externa convexi; articulus quartus minimus tertio excentricè insertus. Pedes ex piceo fusci, setulis paucis instructi. Lamina stigmatica triangularis, angulis late rotundatis, rima stigmatica claviformi. Lamina analis utriusque lateris setae duae in margine postico ferens.

Long. (in satiato) corporis 20 millim.; rostri 1 millim., scuti occipitalis 2-3 millim.; lat. corporis 11 millim., scuti occipitalis 2-25 millim.

Hab. GUATEMALA, Retalhuleu (*Stoll*).

Male unknown. Body of the female ovoid, the skin of the abdomen greyish-brown, marbled with small dark spots and striæ. Occipital plate triangular with rounded angles, on its dorsal surface with black, comparatively large dimples. The general colour of the occipital plate is shining black, with a large amber-yellow spot on its posterior angle, which continues towards the anterior margin as a broad chestnut-brown stripe. On the humeral margin of the scutum the punctiform dimples are wanting. Frontal plate shining, black, with a few punctiform dimples and two oblong frontal holes, truncate behind, angulate in front. Mandibles black at their base, becoming yellowish towards the apex, bearing five hooks on each side, which are arranged two and three on two branches. Palpi shining, dark brown, bearing a few short bristles, abruptly depressed or almost concave on their inner surface, convex externally, the fourth joint inserted excentrically on the third one. Legs dark brown, with some bristles. Stigmatic plate triangular, with rounded angles and a claviform stigmatic fissure. Stigma proper arcuate, with swollen margins.

I accidentally found one specimen of this species among a lot of *A. mistum* collected at Retalhuleu. I dedicate it to my friend Prof. A. Forel, the well-known myrmecologist.

3. *Amblyomma crassipunctatum*, sp. n. (Tab. XIV. figg. 1, 1 a-1 h, d.)

d. Corpus ovatum, postice late rotundatum, depressum, cuticula foveolis punctiformibus sat profundis areis herigatis irregulariter intermixtis instructa. Color obscure rubro-fuscus, maculis albidis secundum margines corporis distributis: macula magna rhomboidea regionem humeralem utriusque lateris occupat, macule quatuor parvæ in interstitiis 3^o, 5^o, 7^o, et 9^o inter scutulos margine postici siti sunt. In margine laterali maculae duæ vel tres adsunt. Scutum frontale supra platum, nitidum, fuscum, lateribus convexe declivibus, sine foris frontalibus, asculis obliquis irregularibus nec non punctis subtilibus impressis atque striis parvis albidis ornata. Rostrum et palpi clarius rubro-fusci. Palporum articulus quartus in tertio apice centraliter insertus. Area mandibularis in utroque latere bamulis tribus in duos ramos distributis. Hamulorum apices colore succineo nitentes. Hamulus secundus primi rami bidentatus. Dentes maxillares simplices, unicolors transparentes vitrei, dense conferti, squamiformes, modice erecti. Lamina stigmatice parva, triangulato-semilunaris, fissura stigmatice parva, claviformi, rima stigmatice arcuata. Valvula analis utraque lateris setis quatuor armata: tribus in margine interno, duabus in margine postico sitis. Fissura genitalis transversa, in area polygonæ sita, labris intrassatis. Pedes rubro-fusci, ad apicem articulorum albedo-annulati, setulis armati.

Long. corp. 7.5 millim., scuti frontalis cum rostro 2 millim.; lat. 5 millim.
♀ latet.

Hab. NICARAGUA, Chontales (Janson).

Body of the male oval, depressed, dark reddish-brown, with many deeply impressed, punctiform dimples, between which are a few callous, smooth, irregularly distributed spots. A comparatively large rhomboid whitish spot adorns the shoulder-region of each side, and on the 3rd, 5th, 7th, and 9th interstices between the furrows of the posterior margin there is a small whitish spot. A few irregular spots of the same colour are seen on the side margin. The frontal plate is flat, shining, with some oblique furrows and impressed dimples; a few whitish stripes interrupt the light chestnut-brown

colour of the plate. The rostrum and palpi are light chestnut-brown; the fourth joint of the palpi is centrally inserted. The mandibular hooks, three in number, are of a shining amber-yellow at the apex; the second hook of the first branch is bidentate. The maxillary teeth are transparent, uniformly white, simple, scaly, moderately erect. The stigmatic plate is small, its shape between triangular and obliquely semilunar; the stigmatic fissure short, claviform, the stigma proper small and arcuate. Each of the anal valvulae bears five bristles. The genital fissure, which is comparatively large, lies in a polygonal cavity and is enclosed by a sort of swollen lip. Legs reddish-brown, whitish at the apex of the joints.

Female unknown.

The figures and description are taken from a dried specimen obtained in Nicaragua by Mr. Janson.

4. *Amblyomma sabanerae*, sp. n. (Tab. XIV. figg. 2, 2 a-2 i, ♀.)

♂ latet.

♀. Corpus ovatum, depressum. Scutum occipitale nitidum, sat dense et haud profunde punctatum, colore ex fusco nigrescente, secundum marginem humeralem utriusque lateris macula lineari irregulari albida, quae in exemplo desiccato diffculter observatur. Scutum frontale nigro-fuscum, foris frontalibus profundius, ovatis, antice angulatum protractis. Abdomen haud nitens, crasse punctatum; superficies dorsalis auleis longitudinalibus latis iniqua. Rostrum et palpi fasci, clariore apicem versus. Palporum articulus quartus in tertio apice excavato centraliter insertus. Arca mandibularis hamulis quique in ramis duos dispositis: hamulus secundus primi rami bidentatus, hamulus primus secundi rami brevioribus. Dentes maxillares acuti, sat dense conferti, in medio apicem versus torulo mucosae incrassati. Lamina stigmatica triangularis, angulis rotundatis, fauce stigmatica irregulariter claviformis, lata; rima stigmatis irregulariter arcuata. Valvula analis utriusque lateris setis quinque armata. Apertum genitalis marginis posteriore rostrato, laterali flexuoso convexo; inter regulas integumentum parietis anterioris aperturam genitalis toruli lineares recti observantur. Pedes clarius fusco-rufi, eorum dentibus duobus respicientibus armati.

Long. corp. 7 millim.; scuti occipitalis 2 millim.; lat. corp. 5 millim.

Hab. GUATEMALA, Retalhuleu (*Stoll*).

Male unknown. Body of the female oval, depressed. Occipital plate shining, punctate, brownish-black, with an irregular whitish stripe on the shoulder-margin, which almost disappears after desiccation. Frontal plate brownish, the two frontal holes deep and oblong. Abdomen not shining, rather coarsely punctate, with some broad longitudinal furrows on its upper surface. Rostrum and palpi brown, lighter towards the apex; the fourth joint of the palpi centrally inserted on the third one, which has its apex excavated. Five mandibular hooks arranged on two branches; the second hook of the first branch bidentate, the first one of the second branch very short and rudimentary. Maxillary teeth acute, with an amber-yellow longitudinal swelling on the middle, their margin double-bordered. The stigmatic plate is triangular with rounded angles; the stigmatic fissure irregularly claviform, comparatively broad, the stigma proper moderately arcuate. Each anal valvula bears five bristles. The posterior margin of the genital orifice is straight, transverse, its side margin convex, arcuate; a

series of short linear swellings between the ripples of the integument of the anterior margin. Legs light brown. The coxæ bear two broad, rounded teeth.

I found two females of this species attached to the throat and tail of a small terrapin, known to the natives by the name of "la Sabanera." The figures and description are taken from specimens preserved in alcohol.

Fam. ORIBATIDÆ.

[Dugès, Rech. sur l'ordre des Acar., 3^e Mém., in Ann. Sci. Nat. sér. 2, i. p. 21, *Oribatei* (1834);
Kramer, Grundzüge zur System. der Milben, in Arch. f. Naturg. xliii. p. 215 (1877).]

Subfam. PTEROGASTERINÆ.

[Michael, British Oribatidæ, pp. 61 & 202 (1884).]

ORIBATA.

Oribate, Latreille, Hist. nat. gén. et partic. des Crust. et des Ins. vii. p. 400 (1804); Gen. Crust. et Ins. i. p. 148 (1806).

1. *Oribata centro-americana*, sp. n. (Tab. XV. figg. 1, 1a-1f.)

Corpus globosum, supra valde convexum, nigrum, leve, nitidum, sine sculptura microscopica ulla. Domo-vertex ab abdomine non separatus, antice sulco transverso ab rostri tecto separatum. Rostrum tectum, valde declivum, triangulare, antice late angulatum, simplex. Lamellæ brevæ, angustæ, apice brevi, acutæ, seta mediocris longitudinis armatæ. Pseudostigmata ab dorsa insipienti non perspicuntur. Organa pseudo-stigmatica longæ, setiformia, apice paululum incrassato. Tectopodia non conspicua. Pedes rufifusci, mediocris longitudinis, reticuli armati, coxæ, femoribus atque tibiis posticis complanatis; tarsus unguibus tribus instructus. Pteromorphæ mediocræ, antice retractæ, late rotundatæ, plerumque corpori appressæ; ita ut a dorsa insipienti difficulter videantur, colore rufo-fusco, semi-pellucidæ. Laminae anales semilunares. Palpi nihil extraordinarii exhibent. Mandibulae crassæ, brevæ, brachia chalarum dentibus quatuor obtusis armatæ. Maxillæ incisuram longitudinalem margini externo vicinam profundam præbent, quæ partem anteriorem marginis externi dentis inter diocernit.

Long. corp. 0.9 millim.; lat. corp. 0.8 millim.

Hab. BRITISH HONDURAS, R. Hondo, R. Sarstoon, Belize (*Blancaneaux*); GUATEMALA, Antigua, Guatemala city (*Stoll*); PANAMA, Volcan de Chiriqui 2500 to 4000 feet (*Champion*).

Body globose, of a short and almost circular shape, broad and rounded behind; colour black, the legs and edges of the pteromorphæ brown; texture polished, shining; without any hairs or minute sculpture on the dorsal surface. Cephalothorax comparatively broad; rostrum simple, broadly pointed; lamellæ small and narrow, short, ending in a short acute point; from the anterior margin of the lamellæ there project two comparatively short bristles on each side. Tectopodia wanting. Pseudostigmata short, difficult to be seen from above, projecting from the angle between the basis of

the cephalothorax and the anterior insertion of the pteromorphæ. Pseudostigmatic organs moderately large, slightly increasing in size towards the ends, a little recurved on the notogaster. No interlamellar hairs conspicuous. Coxæ and femora of the two posterior pairs of legs rather flattened. Notogaster not separated from the dorso-vertex, entirely hairless. Pteromorphæ middle-sized, flexible, semi-transparent, and light-coloured; projecting obliquely forward, and having rounded anterior ends, when seen from the side (from the dorsal aspect their anterior ends seem rather pointed). Genital plates widely separated from the anal plates, occupying a sort of pentagonal area, whilst the anal area is rather circular. Mandibles bearing four blunt teeth on each branch of the chelæ. Anterior margin of the maxilla showing a deep longitudinal fissure near the outer edge.

This species seems to be common and widely distributed throughout Central America. I have found specimens of it in Antigua and in Guatemala city, under stones and on the moist walls of the house-wells (pilas). It has also been collected in British Honduras by M. Blancaneaux, and on the Volcan de Cbiriqui by Mr. Champion. One of the dried specimens from British Honduras which I dissected contained about twenty oval eggs, of 0.3 millim. length.

2. *Oribata rugifrons*, sp. n. (Tab. XV. figg. 2, 2 a-2 d.)

Corpus ovoidum, supra convexiusculum, antice attenuatum, postice late rotundatum, nigrum, nitidum, nudum, supra sculptura microscopica a rugulis longitudinalibus subtilissimis confecta oratum, quæ in cephalothorace magis conspicua, in notogastro paululum obsolete sunt. Cephalothorax ab abdomine solum transversali discretus, valde declivus, conicus, antice late angulatus, simplex. Rostri clypeus a dorso-vertex non discretus. Lamellæ haud conspicuæ. Pseudostigmata recondita; organa pseudo-stigmatica medioarea, in animalculo desiccato setiformia, apice incrassata. Setæ interlamellares abeunt. Testopodia haud conspicua. Pedes mediocres, colore fusco, setulis et unguibus tribus armati. Pteromorphæ longæ, sat angustæ, antice angulatis protractæ, rotundatæ, postice acutiusculæ, mobiles: corpori appressæ pedes omnes obtegent. Abdominis latera ad recipiendâs pteromorphas leviter excavata. Sculptura microscopica pteromorpharum a rugulis radiatim ex insertione setarum fundentibus confecta. Laminæ anales semilunares, genitalibus majores. Laminæ genitales ab analibus valde distinctæ, aream rotundato-pentagonam occupant. Mandibulæ crassæ, ebeharum brachium fixum dentibus 4, brachium mobile dentibus 6 armatum. Palporum articulus secundus incrassatus, quintus in medio dentem unguiformem acutum, curvum gerens. Long. corp. 0.85 millim.; lat. max. 0.6 millim.

Hab. BRITISH HONDURAS, Belize (*Blancaneaux*); GUATEMALA, Retalhén (*Stoll*).

Body ovoid, convex, rounded behind, its anterior end somewhat blunt; black, shining, but not polished, without hairs on its dorsal surface, this latter showing a minute sculpture consisting of closely placed, extremely fine, longitudinal wrinkles, which are more distinct on the cephalothorax than on the notogaster. A transverse furrow separates the notogaster from the cephalothorax, which is very simply built, conical, ending in a somewhat blunt point. Hood of the rostrum forming but one piece with the cephalothorax. Lamellæ wanting. Pseudostigmata apparently hidden, invisible from above; pseudostigmatic organs in the dried specimens of moderate size, setiform, increasing

BOL. CENTR.-AMÉR., Arachn. Acar., September 1891.

g*

towards the end, slightly recurved on the notogaster. Interlamellar hairs and tectopodia apparently wanting. Legs brown, bearing three claws and some bristles. Pteromorphæ long, rather narrow, slightly coloured, brown, semitransparent, their anterior end protracted forward so as to nearly reach the top of the rostrum, rounded, the posterior end somewhat pointed; they are very flexible, and when closed completely hide the legs, the whole body then assuming the shape of a small black seed. The minute texture of the pteromorphæ consists of extremely fine wrinkles, which radiate from their point of insertion. The sides of the anterior part of the abdomen are slightly excavated for the reception of the pteromorphæ. Anal plates larger than the genital plates, occupying a somewhat pentagonal area. Genital plates semilunar. Mandibles short, thick; the fixed branch bears four, the flexible branch five, short blunt teeth. Second joint of the palpi thick; the fifth joint bearing on its middle a claw-like, pointed, slightly curved tooth.

Examples of this species were captured by me in the forests near Retalhuleu, where it lives on dead wood in damp places during the rainy season, in company with *Hoplophora retalteca*.

Note.—*O. rugifrons* being the only true *Oribata* observed by myself in the "tierra caliente" of Retalhuleu, it seems possible that the nymphal form of this genus delineated by me on Tab. XV., figg. 3, 3a-d, belongs to this species. Clusters of hundreds of individuals of these were found crawling on the underside of a piece of dead wood in a cacao-plantation near Retalhuleu on the 3rd of July, 1880. They bear a marked resemblance to those nymphal forms upon which Koch founded his genus *Murcia*, especially to his *M. trimaculata* (C. L. Koch, Deutschl. Crust., Myr. und Ins. Heft iii. p. 136. n. 21).

The following are the notes which I then made on the living animal :—

Body oblong, arched, broadly rounded behind, rather diminishing in size towards the somewhat blunt anterior end. Cephalothorax distinctly separated from the notogaster by a deep furrow, attenuated at its base, conical, large, triangular; palpi projecting on the sides of the apex when seen from above. General colour greyish-white, the dark intestines shining through the smooth and transparent skin. In the angles of the abdomen on each side a globose, yellow, shining corpuscle. The dorsal surface bearing a few long pinnate bristles. On the legs the hairs more numerous, also pinnate, but shorter. First pair of legs the longest, the first and fourth pairs longer than the second and third. Tarsi with one claw only. Mandibles short, large, the fixed branch with four, the flexible branch with three teeth. Maxillæ of each side bearing four differently-sized teeth. The pseudostigmata looking like black eye-like spots. Two long bristles projecting on each side in front of the pseudostigmata. Long. 0.5 millim.

Subfam. *APTEROGASTERINÆ*.

[Michael, British Oribatida, i. p. 64 (1884).]

HOPOPHORA.

[C. L. Koch, Uebers. d. Arachnidenyst. Heft iii. p. 116 (1842).]

1. *Hoplophora retalteca*, sp. n. (Tab. XV. figg. 4, 4 a-4 f.)

Corpus ovatum, convexum, nitidum, glabrum, colore clarius rufo-fusco. Cephalothorax conicus, declivus, mobilis, antice rotundato-acutus. Notogastri latera post cephalothoracem late exserta. Mandibulae crassae, dentibus quatuor in quoque brachio armatae. Dens secundus in brachio fixo extra seriem insertus. Maxillularum margo anterior incisuris minus profundis in dentes tres latus divisa. Pedes boves, tarsus unguibus duobus validis, tertio angustiore armatus.

Long. corp. 1 millim.

Hab. GUATEMALA, Retalhuleu (Stoll).

Body ovoid, arched, shining, without any hairs on its dorsal surface, reddish-brown. Cephalothorax conical, broadly angular towards its anterior ends, the side margins somewhat rounded. Sides of the notogaster behind the cephalothorax slightly excavated. Mandibles large, with four teeth on each branch of the chelae; second tooth out of the row on the blade of the chela. Two longitudinal short fissures divide the maxilla into three broad teeth. Legs short; tarsus bearing three claws, two of which are stronger than the other.

Found during the rainy season on dead wood in damp places in the forests near Retalhuleu.

Note.—I reproduce the drawing and description of the tarsus, as I made them when in Retalhuleu. But still I do not feel quite sure about the correctness of what I saw eleven years ago, as the tarsus of the European *Hoplophora nitens* bears only one claw. Having no preserved specimens of *H. retalteca* at my disposal at the present time I cannot decide the question.

Fam. *NICOLETIELLIDÆ*.

NICOLETIELLA.

Nicoletia, G. Canestrini & F. Fanzago, Intorno agli Acari Italiani, p. 52 (1877).

Nicolettiella, R. Canestrini, Osserv. sulla *Nicolettiella cornuta*, p. 6 (1882).

1. *Nicolettiella neotropica*, sp. n. (Tab. XVI. figg. 1, 1 a-1 c.)

Corpus oblongo-ovatum, convexum, antice attenuatum, margine anteriore bicornute, colore sanguineo. Superficies dorsalis setis sparsis induta. Dorsal cuticula forolis microscopicis rotundis vel hexagonis, quae interstitiis transverse striolatis separatae sunt, regulariter et dense punctulata. Pedes antici ceteris longiores atque crassiores, ceteri graciles; pedes postici secundis atque tertiis paululum longiores. Tarsi primi

et secundi paris unguibus duobus, tertii et quarti paris unguibus tribus armati. Tabercula lateralia valde conspicua, oblonga. Mandibule chelate, crasse, in margine externo sub insertione brachii fixi chelorum dentibus sex brevibus regularibus ornate. Chelorum brachium fixum curvum, acutum, dente valido ante apicem armatum; brachium valde curvum, in medio serrulatum.

Long. corp. 0.5 millim.

Hab. GUATEMALA, Retalhulen (*Stoll*).

Body oblong, convex above, slightly diminishing in size towards the anterior margin, which bears two acute cornicles; blood-red, the legs and cornicles slightly coloured; a few short bristles on the dorsal surface, which shows a minute texture consisting of thickly-set round or hexagonal grooves, separated by transversely striated interstices; on each side between the second and third pairs of legs there projects an oblong tubercle. Anterior pair of legs longer and thicker than the others, which are rather slender; hind legs a little longer than the second and third pairs. The tarsi of the two anterior pairs of legs bearing two, those of the third and fourth pairs three claws. Mandibles short, thick; the fixed branch of the chela slightly curved, pointed, bearing a strong tooth near its apical end; the movable branch strongly curved, its inner edge serrulate from the middle to the base; under the base of the fixed branch the body of the mandible shows a row of six short regular teeth.

Found on one occasion only among fallen leaves in a forest near Retalhulen.

Fam. GAMASIDÆ.

[P. Méguin, Monogr. de la famille des Gamasides, in Journ. de l'anat. et de la physiologie (1876).]

Subfam. UROPODINÆ.

UROPODA.

Uropoda, Latreille, Gen. Crust. et Ins. i. pp. 157, 158 (1806).

1. *Uropoda echinata*, sp. n. (Tab. XVI. figg. 2, 2 a-2 c.)

Corpus supra et infra convexulum, subglobulare, paululum longius quam latius, dorso scutum orale, cornuans, colore brunneo, ab corporis margine membrana albide-grisea late separatum, punctulis densis microscopicis nec non setis brevibus indutum. In membrana conjunctiva marginali series tres oculorum microscopicum ostendit, in quorum quoque seta brevis, spiniformis adest. Tarsus pedum paris primi unguibus caret; ceterorum paris unguis gracilis, acuti. Palpi simplices, breves, crassi. Mandibule longe protrahiles, graciles, chelis angustis, parvis. Chelorum brachium fixum mobili set longius, ad apicem dilatatum, ambo brachia dentibus obtusis indistinctis armata. Scutum ventrale convexulum, foveolis punctatum.

Long. 0.75 millim.

Hab. GUATEMALA, Antigua (*Stoll*).

Body almost globular, a little longer than broad; the dorsal and ventral plates chiti-

nous, reddish-brown, widely and completely separated by a whitish membrane, the latter bearing several rows of small conical protrusions, on each of which is a short thorn-like bristle; the dorsal plate also shows several short bristles and a dense microscopical punctation. The legs are slender, comparatively long; the tarsi of the first pair are without claws, the remaining tarsi with two slender claws. Palpi short, 5-jointed, simple, rather thick. Mandibles, when protracted, long and slender; the fixed branch of the chela is longer than the movable one, both branches being indistinctly and bluntly denticulated. Ventral plate rather convex, dotted with small grooves.

One specimen, found crawling among mould under a hedge in the vicinity of Antigua, Guatemala.

2. *Uropoda inaequipunctata*, sp. n. (Tab. XVI. figg. 3, 3 a-3 d.)

Corpus valde depressum, oblonge oratum, antice in apicem brevem protractum, postice late retundatum, nitidum, colore coccineo. Dorsum microscopio insipienti foretela punctiformes sparsas ostendit, inter quas puncta minora numerosa, marginem versus sece accumulantis aita sunt. Pedes breves, longitudine fere aequales, tarsi primi parie unguibus carentes. Mandibulae longae, graciles; obelae graciles, parvae, eorum brachia irregulariter atque subtiliter denticulata.

Long. 0.5 millim.

Hab. GUATEMALA, Retalhuleu (Stoll).

Body flat, oblong, its anterior border forming a short acute angle, its posterior border broadly rounded, shining, scarlet-red; the dorsal surface shows several punctiform grooves and the space between them is very finely punctured by numerous microscopic points, which are more numerous near the borders. Legs short, almost equal in length; anterior tarsi without claws, the tarsi of the remaining legs with some thorn-like spines. Mandibles long and slender, with small chela, the branches of which are irregularly denticulated.

Found at Retalhuleu as a parasite on a coprophagous beetle (*Pinotus* sp.) called "ronron" by the inhabitants.

3. *Uropoda discus*, sp. n. (Tab. XVII. figg. 4, 4 a-4 c.)

Corpus planum, fere orbiculare, antice leviter protractum, brunnescum, nitidum. Palpi breves, in animalculo sece moventi marginem anteriorem pallidum superantes, eorum articulus quartus tertio longior, ad apicem setis nec non arvis verrucosis, granulatis, circumscriptis nonnullis ornatum. Pedum per primum breve, antevuliforme, unguibus carens, ejus articulus secundus atque tertius dentibus obtusis brevibus irregulariter serratis, articulus ultimus setulis numerosis armatus. Mandibulae longae, tenerae, transverse albidae; obelae corneo-flavidae, parvae, subtiliter denticulatae.

Long. 0.3 millim.

Hab. GUATEMALA, Retalhuleu (Stoll).

Body flat, almost orbicular, its anterior margin slightly protracted, but round. Colour of the body, palpi, and legs brownish-red, shining; mandibles white, transparent, their chela yellowish. Palpi rather short, their last joint showing several granulated

spots and some bristles. Mandibles long and slender, the branches of the chelæ apparently deeply denticulated. Anterior pair of legs rather short, without claws, their second and third joints irregularly denticulated; the posterior pair of legs longer than the others.

I found one example of this species in a decayed chichique-fruit in the woods near Retalhuleu in June 1880. It is nearly allied to the European *U. cassidea*, Herm.

4. *Uropoda centro-americana*, sp. n. (Tab. XVII. figg. 1, 1 a-1 f; 2, 2 a, 2 b, nymph.)

♀ juv. Corpus depressum, oblongo-ovoidum, antice inter secundum et tertium pedum par oblique subtruncatum, ita ut pars anterior scuti dorsalis in sagulum obtusum protrahatur. Superficies dorsalis plana, marginem versus declivis, foveolis numerosis cribata, ex quibus setule breves emaneant, colore brunneo claro (in desiccato animalculo), minus nitente. Pedes omnes anguliferi, sat breves. Articulus ultimus primi paris pedum setis numerosis ornatus, inter quas una longissima juxta ungues sita, ungues in manubrio longo gracili inserti, graciles. Articulus ultimus ceterorum parium pedum setulis sparsis atque spinis duabus juxta ungium pediculus armatus, brevis. Palpi breves, setigeri, in articulo penultimo seta longa, bifida. Palpi maxillares bidentati. Sub apice anteriore scuti dorsalis setulae duae insertae. Mandibulae graciles, chelarum brachia fere longitudine aequalia, dentibus paucis et obtusis in margine interno undulata.

Long. 1 millim.; lat. 0.6 millim.

Hab. NICARAGUA, Chontales (JANSON).

Body (of the young female) depressed, oblong, its anterior margin forming a somewhat obtuse angle; dorsal surface even, marked with numerous grooves, out of each of which a short bristle arises; the colour, in dry specimens, a clear reddish-brown, not very shining. Legs rather short, all of them bearing claws: on the tarsi of the first pair the claws are fixed on a sort of slender petiole and surrounded by numerous long hairs, one of which is very much longer than the others; the tarsi of the second, third, and fourth pairs bear short bristles and two incrassated spines at the base of the petiolus of the claws. Palpi short, bearing numerous long bristles on their last joint; on the inner side of their third joint there is a long, projecting, pinnate hair. Mandibles slender, their chelæ small, blunt at the top, bearing on the inner edges of the brachia four or five obtuse teeth.

(N.B.—Adhering to the posterior abdominal segments of a Guatemalan specimen of the Coleopterous genus *Atractocerus* I found numerous nymphs of a *Uropoda* (figg. 2, 2 a, 2 b) which I am inclined to identify with the above-described *U. centro-americana*, as they have most of the characters in common, except some which may be the result of the differences of the respective stages of development. These differences are as follows:—

Body flat, oblong, not so distinctly angular on its anterior margin, yellowish, transparent, the hairs on the dorsal plate apparently a little longer and less numerous; the incrassated spines of the legs (which in the adult are placed at the base of the claw) attached to the side of the tarsus, one of them on the false joint and the other on the tarsus proper. Length 0.4, breadth 0.25 millim.

Clusters of these nymphal forms attach themselves by viscous threads to the skin of

their host, the threads being strong enough to keep them from falling off, even when dried up or preserved in spirit.

5. *Uropoda piriformis*, sp. n. (Tab. XVII. figg. 3-3 d.)

Corpus ovato-piriforme, subconvexum, antice et postice rotundatum, obscure brunnescens (in arsiocato), pilis brevibus paucis in margine ornatum. Cuticula foveolis numerosis ciliata. Palpi simplices, setigeri, in articulo basali seta crassa armati. Mandibularum chela dentibus numerosis in brachiis ambobus armata. Orificium genitale subtriangulatum.

Long. 1-2 millim.; lat. max. 0-9 millim.

Hab. MEXICO, Jalapa (Höge).

Body longer than broad, somewhat piriform, a little convex on the back, naked, except for a few short scattered hairs on the margin; dorsal and ventral plates rather coarsely punctured. Palpi simple; one strong bristle on the inner side of the basal joint. Anterior margin of the epistome lobed; two strong bristles on the ventral surface of the exterior lobes.

One specimen.

Subfam. GAMASINÆ.

MEGISTHANUS.

Megisthanus, T. Thorell, Descriz. di alc. Aracnidi inferiori dell' Arcipel. Malese, in Ann. Mus. Genova, xviii. p. 48 (1882).

1. *Megisthanus gigantodes*, sp. n. (Tab. XVIII. figg. 1-1 d, ♀; 2-2 g, ♂.)

♂. Corpus longius quam latius, inversa subovatum, antice attenuatum, ita ut subpiriforme appareat, supra modice convexum, nitidum, andum exceptis pilis paucis sparsis ad marginem cephalothoracis anteriorem in utroque latere sitis, suberectis, nec nec setis longis sex vel septem in utroque latere marginis posterioris in series oblique dispositis, æquidistantibus. Color ferrugineo-fuscus, in nonnullis obscurior, in aliis, præcipua in junioribus, clarior. Scutum dorsale in lateribus et postice limbo albedo angusto cinctum; texturam microscopica a areolis polygonis confectam præbet, inter quas foveolæ rariores impressæ sunt. Latitudo maxima post medianâ corporis invenitur. Margo integer, subcompressus, postice late rotundatus, antice truncatus. Venter (in desiccatis et in alcoholo preservatis) in medio convexus, ad latera declivis, leviss; ejus limbus angustissimus, albicans. Scutum sternale angustum, set longum, leve, ad maximam partem ex rufo albicans, margine anteriore truncato, posteriore rotundato, lateribus ob coarctum insertionem sinuatis. In ejus parte anteriore loter coxas secundi paris pedum orificium genitale parvum, circulare situm est; ad marginem posteriorem in utroque latere foveam vallo albedo lato circumdatam videt, ex qua spinula fortis nascitur. Scutum anulo a sternali interstitio albedo angustissimo separatum, obscure ferrugineo-fuscum, margine anteriore excavato, ita ut marginem posteriorem scuti sternalis recipiat, marginibus lateralibus disjunctis, rotundatis, margine posteriore corporis apicem analem attingens; orificium anale parvum, oblonge ovatum, inter scuti medianum atque marginem poticum situm. Pars anterior et latera scuti sternalis nec non scutum anale et scuta lateralia texturam microscopica a scutellis anastomosantibus confectam præbent. Scutum laterale utriusque lateris integrum, postice post coxas quarti pedum paris in angulum prope marginem situm apice rotundatum desinens, a scuto anali interstitio albedo, angusto, marginem versus sese dilatanti separatum. Spinæculum facillime distinguendum, juxta coxam quartam externe situm, oblongum, marginem anteriorem corporis juxta coxam secundi paris trabecula forti, elevata attingens. Pedum par primam gracile, unguibus caræne, antenniforme, longum, setis sparsis, que ad tarsis apicem magis numerosæ apparent, munitum. Secundum et tertium pedum paris fortis, fusiformis, inferius, setis solum munita, que in

femur dorso spinas simulat. Cuticula ad eorum basin in dentes formam paululum protrusa. Femora quarti paris supra tuberculis quatuor equidistantibus munita sunt, ex quibus setae fortes spiniformes nascuntur; subius apicem versus dentes tres fortes gerunt. Cetera articuli setis sparsis muniti, ex quibus una ad apicem articuli quarti, altera ad apicem quinti articuli longiores sunt. Epitoma magnam, triangularem, supra transverse compressam, antice paulo deflexam, apice in angulum scutum paulo producto, eandem texturam microscopice exhibens quam scuta ventralia. Ab animalculo vivo epitoma ad latera nec non aures deorsumque capitis instar meretur. Palpi longi, teneri, setis sparsis quae in apice sexti articuli magis numerose sunt, muniti. Hypotoma in utroque latere dentem apice obtuso ostendit, in medio in linguam longam, angustam, quadrilobatam, setulosam, pellucidam, desinit. Mandibulae fortes, eorum chela longa, angusta, brachiis ambobus denticulatis; in brachio fixo circa 12, in mobili circa 14 dentes observari. Chela brachium fixum in superficie interna scopa pellucida setulosa superatur quae ex chela media nasci videtur; apex brachii mobilis membranula tenuissima, hyalina, convoluta (mappula apud Thorell) ornatur, cujus margo subtilissime serrulatus est. In superficie laterali brachii chela mobilis artusculi tres ramosi nascuntur, quorum duo sub apice brachii siti, et valde inter se approximati sunt, tertius paulo distans, in medio chela fere situs. Ad basem externam brachii mobilis fasciculus ex spinulis hyalinis confectus diffillime observatur. Dentes labiales longi, curvi, integri.

Long. corp. cum epitom. 3.5 millim., sine epitom. 3 millim.; lat. max. 2-2.25 millim.

♀. Corpus ovalum, antice truncatum, supra convexulum. Scutum dorsuale integrum, modice nitidum ob foveolas impressas, ex quibus setulae breves, subrectae nascuntur, oblique retrospicientes. Limbo albido ad latera et postice a scuta ventralibus separatur. Color late brunneus. In utroque latere prope marginem scoti dorsalis series setarum longarum inserta est. Textura microscopica e sulculis anastomosantibus confecta, qui scutum totum areolis oblongis irregulariter reticulatis, sicut in maris scutis ventralibus. Venter in medio convexus, lateribus in coxarum regione depresso-convexis. Scutum sternale angustum, antice et postice rectum, lateribus ob coxarum insertionem sinuatis, a scutis lateralibus interstitio angusto, a scuto anali interstitio lato albicanti discretum. Inter coxas secundi atque tertii pedum paris orificium genitalis situm est, magnum, subpentagonum, margine postico incrassato. Scutum anale margine antico recto, lateribus atque margine postico rotundatis, orificio anale in medio situm. Scuta lateralia coxiferis ab anali interstitio albedo lato separata, postice angulatis protrata, apice rotundata. Epitoma, palpi et pedes sicut in maris. Hypotoma in medio in linguam quadrilobatam hyalinam productum, dentibus lateralibus, qui in mare adnascuntur, carens; sulculo laterali sinuato, qui palporum socalem a labii tranco separat. Palpi labiales longi, graciles, apice scuto, curvo, margine interno dente uno seu duobus. Mandibulae sicut in maris, excepto brachio mobili chela, qui numerum minorum dentium praebet.

Long. cum epitom. 2.5 millim.; lat. max. 1.25 millim.

Hab. GUATEMALA, Acuituno, Guatemala city, Zapote (*Champion*), Retalhulen (*Stoll*). In regione "Cholhuitz," hanc speciem in coleoptero-pectinicorni "*Proculo gorgi*" denominato parasitam inveni.

♂. Body longer than broad, rounded behind, truncate at its anterior margin, much depressed, smooth and shining; a few hairs are placed near the anterior margin, and a row of six or seven obliquely inserted long bristles adorn the hind margin on each side; colour light reddish-brown, in some individuals darker than in others; the dorsal plate is separated from the ventral pieces by a whitish border on its side- and hind-margins, and shows a minute sculpture consisting of small, irregularly polygonal areas with a few impressed points between them; ventral surface convex in the middle, depressed at the sides, smooth, its border whitish and very narrow. Sternal plate narrow; rather long, smooth; its anterior margin truncate, its hind margin rounded. Genital orifice situated between the coxae of the 2nd and 3rd pairs of legs, small, circular. The sternal plate has two grooves at the hind margin, one on each side, out

of each of which a strong hair arises; the grooves are encircled by an apparently elevated broad whitish border, the border showing a microscopical granulation.

These characters are peculiar to the male sex of the genus *Megiothanus*, and are wanting in the female; the colour of the sternal plate itself is much lighter than that of the anal plate, and it is somewhat irregularly distributed—darker on the anterior and lateral parts, lighter on the middle; the anal orifice is situated in the middle of the anal plate. The side-plates extend beyond the hind coxæ, forming a rounded angle at their end. The anterior legs are antenniform, long, and without claws; at the top of the tarsi there are some densely-set tactile hairs. The second and third pairs of legs are strong, somewhat fusiform, bearing on the back of the femora some spine-like hairs. The posterior femora have on the back an equidistant row of four spine-like hairs and on their lower surface three strong, short, blunt teeth. The remaining joints of the legs are irregularly set with hairs. The epistome forms a large triangular hood, which is convex above and ends in a short acute angle; during life this piece can be moved upwards and laterally; it hides the oral parts but imperfectly, as it is surpassed in length by the palpi and, sometimes, by the upper part of the mandibles. The palpi are long and simple, sparingly set with hairs, which are more numerous at the top of the fourth and fifth joints. The margin of the hypostome bears a blunt tooth on each side, and its middle projects into a transparent quadrilobate tongue-like piece. The mandibles are strong, their chelæ rather narrow, bearing numerous irregular teeth on the inner edge; on the inner surface of the fixed branch of the chela a transparent narrow brush projects a little over the top of the mandible; the movable branch seems to be enveloped at the top in a small, transparent, irregularly-folded membrane, the border of which is serrulated; on the lateral surface of the movable branch there project three transparent appendages, each of which bears some finger-like ramifications, and at the base there is a transparent cluster of small spinules.

♀. Body ovate, its anterior margin truncate, slightly convex on the dorsum; dorsal plate but moderately shining, rather coarsely punctured, a short semi-erect bristle arising from each puncture; colour reddish-brown, the border of the dorsum whitish. On each side of the dorsal plate, along its lateral margin, a row of long bristles is inserted. The microscopical texture is similar to that of the ventral pieces and hood of the male; it consists of a network of fine anastomosing furrows, which leave long, narrow, irregular areas between them. Ventral surface convex in the middle, depressed and concave at the sides. The whitish interstice between the sternal and lateral plates is narrow, that between the sternal and anal plates being broader. The genital orifice, which is comparatively large and pentagonal, lies between the 2nd and 3rd pairs of coxæ. The anterior margin of the anal plate is straight, its side-margins being rounded; the anal orifice is situated in the middle of the plate. The lateral plates project into an obtuse angle behind the hind coxæ. Epistome and palpi as in the male. The hypostome bears no lateral tooth; the tongue-like piece is formed as in

the male. The movable branch of the chelæ has fewer teeth than in the male; but the fixed branch and appendages are similarly formed.

This is the largest Gamasid known as yet from Central America. In the Cholutz district near Retalhuleu I found it parasitical upon a very large pectinicorn-beetle, *Proculus goryi*, Guér.; but it cannot be confined to this species only, as the genus *Proculus* is not known to occur in the localities where Mr. Champion obtained his specimens of *M. gigantodes*. It is perhaps parasitic on other Passalidae.

2. **Megisthanus armiger.** (Tab. XIX. figg. 1, 1 a-e, ♂.)

Megisthanus (sic) *armiger*, Berlese, Bull. Soc. Ent. Ital. xi. p. 204, t. 9. figg. 1, 1 a (1888).

Hab. MEXICO, Jalapa (Höge).—PARAGUAY, Rio Apa.

Among the dried specimens of Gamasidæ received from Messrs. Godman and Salvin there is one male *Megisthanus*, collected by Herr Höge at Jalapa, which I believe to belong to the above-named species, as it coincides with the description in all its essential characters, as well as in the measurements, though it differs from it in some minor details.

The differences are as follows:—

<i>Specimen (♂) from Jalapa (Mexico).</i>	<i>Specimen (♂) from Rio Apa (Paraguay), as described by Mr. Berlese.</i>
Scuti dorsalis superficies pilis brevibus e foroleis protuberantibus nec non punctulis impressis ornata. In dorsi medio area oblonge tri- angularis sulculo lineari circumscripta adest.	Scutum dorsuale et marginæ pilis cartulis vestita. (Nothing more mentioned.)
Ante scutum sternale scutulum prosternale, parvum, transversum, medio longitudinaliter partitum sicut est.	(Not mentioned.)
Scutum sternale postice rotundatum, anterior ex- caratum. Scutum ventrale est ab anali (trans- verso quadrilatero) discretum.	 rotundatum, in medio tamen sinuatum, anterior truncatum. Scutum ventrale parum ab anali (triangulari, vertice infero) discretum.
Margo metapodiorum internus excarvatus.	(Straight in Berlese's drawing.)
Segta ventralia omnia setulis brevibus vestita.	Metapodis et scutum ventrale setis vestita, cetera nuda.
Femora tertii paris inferne calcaribus tribus fere obsoletis, quarti paris calcaribus quatuor bre- vibus obtusis aucta.	Femora tertii et quarti paris inferne calcaribus tribus validis aucta.
Corniculi labiales curvi.	Corniculi labiales recti.
Trochanter secundi paris superne inermis.	Trochanter secundi paris superne spina valida armatus.
Cuticula in articulo quarto et quinto secundi, tertii et quarti pedum paris ante apicem in dentem squamiformem subtus sublevata.	(Not mentioned.)
Articulus palporum secundus dento brevi ante api- cem subtus armatus.	(Not mentioned.)

CELENOPSIS.

Celenopsis, Berlese, Acari, Myr., et Scorp. Ital. fasc. xxxi. no. 5 (1886).

Diplogynium, Canestrini, Atti Soc. Ven.-Trent. xi. p. 101 (1888).

1. *Celenopsis uropodoides*, sp. n. (Tab. XVI. figg. 4-4 d, and Tab. XIX. figg. 3, 3 a, b, ♀.)

♂ latet.

♀. Corpus oblongo-ovatum, depressum, supra minime convexum, postice indistincte acuminatum. Scutum dorsale limbo angustissimo, subnitidum, setulis brevibus in superficie et in margine suberectis munitum. Color ladius. Sculptura microscopica e tubulis anastomosantibus confecta, qui areolas oblongas irregulares circumambulant. In harum areolarum medio puncta seriatim disposita impressa sunt. Palpi et pedes marginem corporis superant. Superficies ventralis depressa, eadem structuram microscopica sicut dorsalis præbens. Scutum ventrale magnum, postice acuminatum, marginem posteriorem corporis non attingens, orificium anale prope apicem ferens. Coxæ utrinque lateris approximatae. Inter coxas secundi etque tertii pedum paria orificium genitale (a clarissimo hujus generis auctore "epigynium" appellatum) in parte anteriore scuti ventralis situm est, magnum, subpentagonum, antice latum, angulo postico rotundatum. In epigynio distinguenda sunt:—(1) crista epigynica jugiformis, lateribus deflexis, transversa, antice arcuata; (2) valva vulvæ utriusque lateris aliformis, interne atque antice rotundata, ad angulum internum anticum et prope marginem externi medium seta mucronata; (3) scutum parvum trigonum a margini postico epigynii inter valvas utrinque lateris paullo prodians; (4) fulcræ valvæ claviformis, obliques, arcuatae. Ante epigynium scutum sternale hexagonum situm est, cuius margo anterior excavatus, lateralis in angulum rotundatum productus est. Ante angulum et prope marginem posteriorem in utroque latere setula inserta est. Scuta lateralia angusta, longa, prope marginem corporis posteriorem rime triangulari angusta, albide separata. Spiraculum parvum, prope coxas posticas externe situm. Pedes primi paria graciles, eorum tarsus unguibus curvus, longus, setis numerosis ad apicem tacitibus, sicut articulus palporum extremus, munitus. Pedes ceteri inermes, crassiores, setulis muniti. Palpi inermes, set longi. Epistoma trigonum, apice in mucronem acutum desinens. Mandibularum chela brachiis gracilibus, denticulatis; dentes brachii mobiles inaequales; ad apicem parvuli, conferti, infra inter se distantes, majores, pauci; in brachio fixo subaequales, apicem versus magis conferti ac breviores. Hypostoma in linguam antice rotunde truncatam desinens, ex qua processus duo longi setiformes nascuntur. Corniculi labiales basi lata in cornu extensus curvum acutum pellucidum desinunt. Brachium mobile membrana pellucida marginibus sinuatis longa imperfecte obtectum. Apparatus cibarius totus difficillime aruenda.

Long. 0.75 millim.; lat. 0.50 millim.

Hab. BRITISH HONDURAS, R. Hondo (*Blancaneaux*).

Male unknown. Body of the female ovate, depressed, moderately shining, bearing some short suberect hairs on all parts and showing a microscopic texture, consisting of a network of very fine irregular anastomosing furrows, which leave long irregular spaces between them, each space having a series of impressed points. The ventral plate is large and acute behind, and does not reach the end of the body. Anal orifice small, oblong, situated in the posterior angle of the ventral plate. The space between the two rows of coxæ rather narrow. The genital orifice (called "epigynium" by Mr. Berlese) is situated in the anterior portion of the ventral plate; it is large, somewhat pentagonal in shape, broad at its anterior margin, rounded behind. The sternal plate is hexagonal, excavate at its anterior margin, the lateral margin forming a rounded angle. The lateral pieces are long and narrow, separated from the ventral piece by a

narrow interspace; at the end of the body there is a small triangular space between them. The spiracle is small, distinctly visible, and situated outside the hind coxæ. The legs are without spines or teeth, and bear only some hairs and bristles, which at the top of the first pair of tarsi (and on the palpi) are long and more thickly set than on the other joints. The epistome is triangular and protrudes from under the anterior margin of the dorsal plate as an acute point; it imperfectly covers the mandibles, the branches of which are irregularly denticulated.

Several specimens of this species, which is closely allied to the South-American *C. subincisa*, Berlese, were collected at Rio Hondo by M. Blaneauaux.

2. *Celénopsis megisthanoides*, sp. n. (Tab. XIX. fig. 4, and Tab. XX. figg. 1, 1 a, b, 2.)

♂ latet.

2. Corpus oratum, depressum, antice et postice declivum, postice subacuminatum, antice, propter epistomatē apicem, in angulum scutum declivens, lateribus in regione humerali sinuatis. Color clare ladius. Scutum dorsuale integrum, antice truncatum, ejus margo in lateribus et in margine posteriore; limbo angusto albedo ab acutis ventralibus separatim, setis aut longis sparsis in superficie ornatum. Scutum stercorale parvum, transversum, inter coxas primi paris altum, antice recte truncatum, lateribus et margine postice sinuatis. Scutum ventrale magnam, marginem posticum fere stringens, in eius parte anteriore vulva, rimam bifidam simulans, sita est. Pedes omnes setulis sparsis muniti fere inermes, exceptis femoribus quarti paris quæ calcareibus tribus valde obtusis et seta valida basi approximata armata sunt, nec non tarsis secundi paris qui ad apicem juxta unguitum pedunculum dentem curvum exhibet. Pedes antici longi, graciles, ceteri crassiores, in eorum femorum dorso setæ duæ validæ adsunt. Epistoma trigonum, declivum, antice acutum, ad eius basin in margine scuti dorsalis seta fortis in utroque latere adest. Corniculi labiales longe, ensiformes, extus recurvi, dentibus carentes. Inter eorum basin atque hypostoma processus spatuliformis acutus transparentes adest. Hypostoma in linguam longam, angustam, tripartitam declivens. Chelarum brachia longa, angusta, irregulariter denticulata, brachium mobile dentem fortissem latum prope basin exhibens. Palpi setigeri, inermes.

Long. 1.25 millim.; lat. 0.75 millim.

Hab. PANAMA, Bugaba (*Champion*).

Male unknown. Body of the female ovate, depressed, somewhat acuminate behind, the lateral margin in the region of the shoulders somewhat sinuated; anterior border of the dorsal plate truncate, bearing a strong bristle on each side over the basis of the epistome; the lateral and hind margins of the dorsal plate show a narrow whitish border; the dorsal surface bears some comparatively long but thinly-set hairs. The sternal piece is small, its anterior margin truncate, the lateral and hind margins sinuated; it occupies the space between the first and second pairs of coxæ. The ventral plate is large and almost reaches the hind margin of the body. The vulva is situated in the anterior portion of the ventral piece; it forms a sort of bifid fissure. The fore legs are long and slender, the other legs shorter and thicker; all of them are without spines or teeth, except the second pair, which bear a small curved tooth at their top, and the femora of the hind pair, which underneath bear a row of three somewhat blunt teeth; between the base of the hind femora and the first of these teeth there is a

strong bristle. The labial horns are long and slightly curved, in the shape of the letter **s** reversed; between their bases and the hypostome there is another process, the top of which forms an acutely-pointed transparent foliole or spatula; the hypostome ends in a long and narrow tongue-like process.

Two specimens.

PACHYLELAPS.

Pachylaps, Berlese, Acari, Myr., et Scorp. Ital. fasc. li. no. 10 (1888).

1. *Pachylaps hæros*.

Pachylaps hæros, Berlese, Bull. Soc. Ent. Ital. xx. p. 106, t. 8. figg. 2, 2 a-e.

Var. *mexicanus*. (Tab. XIX. figg. 2, 2 a-e.)

♂. Corpus late ovatum, supra convexum, antice subangulatum, postice rotundatum, antice et postice deditum. Color scutorum, rostri pedumque huius, membranarum conjunctivarum albus. Scutum dorsale integrum, fere nudum, dense reticulatum, setis sparsis in parte anteriore et in marginibus munitis, a margine postico membrana conjunctiva setulis numerosis gerente separatum. Pedes primi parvis gracilibus, articulis plus minusve cylindricis, setis sparsis munitis, tarsis unguiferis. Pedes secundi parvis crassiusculis, dentibus seu calcaribus ut sequitur armati; dentes breves duo in coxae apice adsum, trochanter est isomnis, femur calcaris fusco, recto, indistincte tridentato armatum, genu atque tibia tuberculum parvum bigibbosum gerunt, tervus ante apicem dentem fortem obtusum præbet. Coxæ terti parvis dentibus duobus, uno centrali, altero marginali, brevibus munitis, trochanteres ad apicem dentes duos juxtapositos dempra, tertium infra ferunt, ceteri articuli pilis exceptis inermes. Coxæ quarti parvis dentibus duobus brevibus sicut in tertio pari munitis, trochanteres ad apicem dempra dentes duos subacutos juxtapositos, infra dentem unum obtusum gerunt, femur ad basem et in apice dente armatum, ceteri articuli, tarsis inermibus exceptis, dentibus duobus armati. Scutum sternale cum ventrali concretum reticulatum, postice truncato-rotundatum, marginem posticum non attingens, ab anali separatum. Scuta metapodica parva, trigona angulis rotundatis. Scutum anale parvum, trigonum. Orificium genitale parvum in margine anteriore scuti ventralis situm. Inter scutum ventrale et hypostoma lobulus parvus linguiformis, setulis duabus ad apicem munitis mobilis insertus est. Chela mandibularum modica, dentibus paucis armata, in brachio mobili processu longus ad chela basin in spiram phories convolutus insertus est. Corniculi labiales distincte biarticulati, eorum articulus postremus ante apicem dentem parvum gerit. Palpi inermes, excepto articulo primo inferne calcar robustum ferente.

♀. Statura et coloris maris; a mare precipue differt pedum pari secundo ceteris haud crassiore, inermi, nec non quarto pari calcaribus carente, inermi.

Long. 2 millim.; lat. 1.5 millim.

Hab. MEXICO, San Andres Tuxtla (*Salté*).

Body of the male broadly ovate, slightly convex, its anterior border forming a short acute angle; colour of the hard plates, the rostrum, and legs light reddish-brown, the connecting membranes whitish; the dorsal plate almost hairless, bearing only a few bristles on its anterior and lateral borders; the bristles are longer and more thickly set on the whitish membrane which connects the dorsal plate to the ventral pieces; the dorsal plate shows a minute reticulation, consisting of a network of small polygonal areæ; almost on the middle of the dorsal surface there are two slightly impressed grooves. The first pair of legs are slender, their joints almost cylindrical, bearing some

hairs, but without spines or teeth. The second pair of legs are very thick, the femur especially, the latter bearing a strong, straight, slightly denticulated spine on the middle of its under surface; the genu and tibia each bear a small tubercle, and the tarsus is armed with a broad laminar tooth at some distance from the tip. The third pair of legs are comparatively slender, without spurs, except a short one on the upperside of the trochanter at the tip. The fourth pair of legs are longer than the third, their trochanters bearing two teeth on the upper surface at the apex and a short tubercle below; the femora bear underneath a tooth at their base and one at their apex; the fourth joint has a blunt tooth and a short spine at its tip. The sternal and ventral plates form but one piece of an irregular shape, which protrudes but little behind the fourth pair of coxæ and ends at some distance from the anal plate; this piece has its side-margins excavated for the reception of the coxæ, its posterior portion being widened laterally and truncate behind; the anal piece is small, triangular; the lateral pieces are large and bear the distinctly visible spiracle; the metapodial piece is small, of a triangular or rather oviform shape. The palpi are unarmed, except their first joint, which bears a strong spur on the inner side. The labial horns are distinctly two-jointed, and have a tongue-like process between them. The chelæ bear a few strongly-marked teeth at their edges; from the centre of the movable branch a long process issues, the end of which is rolled up somewhat spirally.

The female differs from the male in the following characters:—All the legs are slender and, with the exception of the apical teeth of the trochanters, do not bear any spurs or teeth; the sternal plate is separated from the ventral piece; there are no metapodial plates; and the dimensions of the body are a trifle smaller.

Two specimens, one male and one female, of this large Gamasid were collected by M. A. Sallé at Tuxtla. Their affinity to *P. hæros*, Berlese, from Brazil, is so remarkable that I can only treat them as a variety of that species. The Mexican specimen (♂) before me differs from Berlese's description in the following characteristics:—

P. hæros, typ. ♂ (as described by Berlese).

Spur of the femur of the second pair thick, curved, with two teeth at the tip; spur of the genu of the second pair comparatively elevated, distinctly visible; tarsus of the second pair without a tooth at some distance from the apex; genu of the fourth pair bearing but one tooth (not mentioned in the description, but present in the drawing).

Var. *mximimus*, ♂.

Spur of the femur of the second pair straight, denticulate at its margins; spur of the genu of the second pair short, rudimentary, forming only a tubercle; tarsus of the second pair bearing a distinctly visible compressed tooth at some distance from the apex; genu of the fourth pair bearing two teeth—one pointed, spined (like the lateral tooth), and one blunt, on the ventral surface of the apex.

HOLOSTASPIS.

Holostaspis, Kolenati, Wien. ent. Monatschr. ii. p. 87, t. 1. figg. 1, 2 (1856); Canestrini, Prospetto dell' Acarofauna Italiana, p. 55 (1885).

1. *Holostaspis marginatus*. (Tab. XX. figg. 2, 2 a-e.)

Acarus marginatus, Hermann, Mémoire Aptérologique, p. 76, t. 6. fig. 6.

Holostaspis marginatus, var. *americanus*, Berlese, Bull. Soc. Ent. Ital. xi. p. 195.

Hab. GUATEMALA, Retalhuleu (*Stoll*); NICARAGUA, Chontales (*Janson*).—SOUTH AMERICA, Brazil, La Plata, Paraguay.

Two specimens of this classical *Gamasus* were collected by Mr. Janson at Chontales. I have frequently met with it at Retalhuleu, where its various larval and nymphal stages live as parasites on the common "ronron" or dung-beetle (*Pinotus* sp.), together with the nymphs and larvæ of *Gamasus fucorum*, De Geer (*coleopratorum*, auct.).

Subfam. *DERMANYSSINÆ*.

PTEROPTUS.

Pteroptus, L. Dufour, Ann. Sciences Nat. xvi. p. 98 (1832).

The bats of Central America are, as well as those of Europe, infested by parasitical mites of the genus *Pteroptus*, which attach themselves to the smooth skin of the wings and the axillary cavity. But as my drawings are too incomplete to allow the identification of the species, I do not reproduce them here, but merely state that this subfamily is represented in Guatemala.

Suborder II. ACARINA-ATRACHEATA, Kramer.

[Kramer, Grundsätze zur Systematik der Milben, in Arch. für Naturg. xliii. p. 218 (1877).]

Fam. SARCOPTIDÆ.

[Michael, British Oribatidæ, p. 50 (1884).]

Subfam. TYROGLYPHINÆ.

[Trouessart, Les Sarcopt. plinicoles, 1^{re} part. p. 6 (1885).]

In Guatemala I have occasionally observed the presence of free-living *Tyroglyphi*, as well as of hypopial forms attached to the body of several species of flies (Muscidæ), but I have not taken exact notes as to any of them.

Subfam. ANALGESINÆ.

[Trouessart, Les Sarcopt. plumicoles, 1^{re} part. p. 5 (1883).]

MEGNINIA.

Megninia, Berlese, Acari, Myz., et Scorp. Ital. fasc. iv. (1883).1. *Megninia pteroglossorum*, sp. n. (Tab. XXI. figg. 5, 5 a, b.)

♂. Corpus subrotatum, colore ex albido-griseo, nitidulum, convexulum. Pedem paria anteriora inter se aequalia, brevia. Pedes tertii incrassati, longi, abdominis marginem posteriorem superantes, articulo ultimo acuto, unguliferi, caruncula adhaerens indurata, ceteris articulis cylindricis. Pedes quarti breves, tenues, in articulo ultimo sagues duo ferentes, abdominis marginem vix attingentes. Appendices anales lamelliformes, acuti, margine interiore integro, exteriori incisuris duabus sat profundis sinuato. Pars posterior appendiculis utriusque lateris setas duas longas ferens, seta brevior in lobo inter incisuras confecta; longissima juxta basin appendiculae externae sita. Disci adhaerivi prope marginem posticum ante appendiculas in ipso abdomine siti, magni. Epimera primi pedum paria in linea mediana ad lamina sternalem formandam sese conjungunt.

Long. corp. circa 0.5 millim.

♀ *nympha*. Corpore oblongo ovato, antice latiore, marginem posticum versus attenuatum, postice rotundatum. Pedes breves, corporis margini inserti, sparse setigeri. Pedes anteriores crassiores, breviores, posteriores tenuiores, longiores. Epimera primi paria in linea mediana sese tanguntia, secundi paria brevia, libera. Marg. posterior abdominis integer, setis quatuor armatus, internis brevioribus, externis longioribus.

Long. corp. circa 0.5 millim.

Larva hexapoda corpore oblongo, margine integro, super pedum insertionem angulariter protracto.

Hab. GUATEMALA, Retalhuleu (*Stoll*). In remigum vexillis *Pteroglossi torquati* parasita gregatim vitam degit.

Body of the male ovoid, greyish-white, transparent, shining, convex. Anterior pairs of legs of equal length, short. Third pair of legs very thick, long, protruding over the hind margin of the body, their last joint pointed, bearing claws, the other joints cylindrical. Fourth pair of legs short, slender, with two teeth on their tarsus, hardly reaching the end of the body. The anal appendices are pointed, lamelliform, their inner margin straight, their outer margin sinuated, with two profound incisures; the posterior portion of the appendices bears two long bristles, and there is a short bristle on the lobe between the two incisures; the adhesive discs are large and situated near the posterior margin of the body before the appendices. The epimera of the fore legs of both sides are united into a common sternal plate.

Body of the female (nymph) oblong, somewhat thinner towards the posterior end, rounded behind. The legs are short and inserted at the margin of the body; the two first pairs are a little thicker and shorter than the hind pairs. The epimera are brown and have the appearance of narrow hard bands; those of the fore legs of each side reach the median line, those of the second pair remain separated. The hind margin of the body bears four bristles; in young specimens it is a little excavated.

The hexapod larva is oblong, the side-margin forming a sort of angle over the insertion of the legs.

This species lives gregariously as a parasite in the wing-feathers of the "Cucharon" (*Pteroglossus torquatus*) in the tierra caliente of Retalhuleu (Guatemala).

PTEROLICHUS.

Pterolichus, Ch. Robin et Mégnin, "Mém. sur les Sarcophtides plumicoles," Journ. de l'Anat. et de la Physiologie, 1877, p. 393.

1. *Pterolichus momotorum*, sp. n. (Tab. XXI. figg. 1, 1 a, b, c; 2, 2 a, 2.)

c. Corpus convexum, translucidum, ex albidis griseum; marginum circumferentia irregulari inter coxas secundi et tertii paria latus utriusque corporis ad lobi instar protrusum, lobulam anteriorem minorem atque posteriorem majorem simulans. A basi tertii paria pedum usque ad basin appendicum analem corporis crassitudine dimittit. Appendices anales magnae, basi lata, margine externo integro, convexo, margine interno irregulari, in medio rotunde excavato, sinuato. Appendiculae apices sat obtusos, setis duabus longis nec non spinis tribus brevibus munitae. Disci adhaerenti in appendicularum basi siti. Pedes omnes inter se aequales, eorum tarsus caruncula magna in pedunculo tenui inserta instructus. In dorsi medio super coxas secundi pedum paria setae quatuor, dum externae longae, aliae sunt. Spina valida recta in latere corporis ante coxas tertiorum pedum. Palpi biarticulati. Mandibulae breves, crassae, eorum chela dentibus paucis brevibus armatae. In facie ventrali acutum sternale parvum, inter coxas pedum anteriorum, atque acutum ventrale, ab illo interstitio clariore separatim, oblongum, distinguuntur. Epimera angusta, linearia, colore branneo. Epimera primi et secundi paria oblique versus lineam medianam tangentia, sed sese non attingentia, apicibus lato separatim. In ventris medio trabecula semicircularis ante orificium genitale sita.

Long. corp. circiter 0.5 millim.

2. Corpus majus atque longius quam in mari, lateribus ad lobi instar protrusum, post coxas quarti paria cylindrica, lateribus parallelis, apice rotundato, in medio rotunde excavato, juxta excavationem in utroque latere setae duae longae et spinae breves duae adsunt. Epimera, setae dorsi, palpi, mandibulae atque pedes ut in mari.

Long. corp. circiter 0.5-0.75 millim.

Hab. GUATEMALA, Retalhuleu (*Stoll*), in vexillis remigum *Momotus lessoni* gregatim.

Body of the male convex, transparent, of a milky-whitish colour; the side-border is irregular and projects between the second and third pairs of legs in the form of a bipartite lobe; from the base of the third pair of legs the abdomen diminishes in size. The anal appendices are large, broad at their bases, convex externally, their inner border irregularly sinuated, with a round excavation in its middle; they are rather blunt at their tip and each bears two long bristles and three short spines; the adhesive discs are situated at their base. On the anterior portion of the back four bristles are inserted, and the side-margin bears, at a short distance from the third pair of coxae, a straight short spine. The legs are of about equal length and shape, each tarsus bearing a large caruncle. The palpi are two-jointed, and the mandibles are short and thick, their chela armed with a few short teeth. The ventral surface is divided by a clearer interstice into a smaller sternal piece, occupying the space between the anterior coxae, and a larger and oblong ventral piece. The epimera form narrow, pointed stripes of a brownish colour, which are directed from the side-margin towards the median line of the body.

Biol. Centr.-Amer., Arch. Acar., January 1893.

G*

The female differs from the male by its larger size and the more oblong shape of the body, the abdomen being cylindrical from the hind legs, and by the want of the anal appendices. The abdomen is simply rounded at the apex, the hind margin bearing a small round excision in its middle. The legs, the dorsal hairs and lateral spines, the epimera, the palpi, and mandibles are as in the male. A chitinated stripe of brownish colour and semicircular shape is situated transversely before the genital orifice.

This species lives gregariously in the wing-feathers of the "Pajaro bobo" (*Momotus lessonii*). In running, the female takes the lead and draws the smaller male after her. In the act of copulation the female places her abdomen above the anal appendices of the male. (In figg. 1, 2 this connection already appears loosened a little.) I observed this species in November 1880.

P. momotorus bears, by the form of its anal appendices, a remarkable affinity to *P. phylloproctus*, Trouess. et Mégn., which inhabits the *Halictus leucogaster* of the Indian and Chinese seas. It seems to be even more closely allied to this species than to *P. biemarginatus* and *P. ramphastinus*, Trouess. et Mégn., which are parasites of various South-American birds.

PROCTOPHYLLODES.

Proctophylloides, Robin et Mégnin, Mém. sur les Sarcophtes plumicoles, p. 629 (1877).

1. *Proctophylloides sialiarum*, sp. n. (Tab. XXI. figg. 3, ♂; 4, ♀ a-c, 2.)

♂. Corpus minimum, albidum, translucidum, circiter quarta parte minus quam in femina, sat longum, penitulum angustius quam in femina, pedes anteriores atque posteriores interstitio longu segregati. Omnes pedes inter sese fere aequales, breves. Maxima latitudo corporis ante coxas tertii paris. Ante eam spina recta prope marginem inferi sita est. Post pedes tertii paris corpus crassitudinem dimittit. Appendices anales sat breves, carnosae, conicae, separate incisura profunda, in quavis appendice 3 setae diverse longitudinis adnatae. Disci adhaerivi in appendicum basi prope marginem inferum inferi siti. Mandibulae breves, chelorum brachia longa, angusta, et ita curva ut clausa apicibus solum sese attingant.

Long. circiter 0.25-0.5 millim.

♀ *adulte*. Corpus majus quam in mare. In abdominis apice furca carnosae adnatae, quae infera prope marginem convexum externum in utroque latere apicem lanceolatum sat longum fert. Furca apex cuiusque lateris in appendicem longam, sinuatam acutissimam desinit. Epimera primi paris utriusque lateris in media regione sternali sese conjungunt, secundi paris angulariter retrosum fructa late discreta remanent. In corporis latere, paululum post medium interstitium inter pedes secundi et tertii paris seta sat longa inserta est. In ventre medium trabecula transversa arcuata seu semicircularis adnata, quae in utroque latere brachio longo analem coxae tertiae attingit. Mandibulae ut in mari.

Long. circiter 0.5 millim.

♀ *nympha*. Abdomen in appendicem conicam, carnosam, incisura bipartitam desinit, quae in superficie externa utriusque lateris apicem lanceolatum sat longum et in apice setam longam gerit.

Larva hexapoda minima, jam rudimenta appendicum analium adultae praebet.

Orum longum, fusiforme, translucidum, sat magnum, ad cribri instar perforatum.

HAB. GUATEMALA, Retalhulen (*Stoll*), in remigibus *Sialia sialis* gregatim.

Male very small, long and narrow, of a whitish, somewhat transparent colour, about one-fourth smaller than the female. The two anterior pairs of legs are separated by a

wide interstice from the third pair. The body is broadest at some distance in front of the third pair of coxæ; from the third pair of legs it diminishes in size, and ends in two fleshy appendices, of a short conical shape. The appendices are separated by a deep incision; each of them bears three bristles; the adhesive discs are situated near the inner margin and at the base of the anal lobes. The mandibles are short; the branches of the chelæ are long, slender, and curved, and when closed they touch each other with the points only.

The adult female is a little larger than the male. Its abdomen terminates in a fleshy fork, each point of which bears a lancet-like spine underneath and ends in a narrow, acutely-pointed, somewhat curved blade. The apices of the first pair of epimera of each side touch each other in the sternal region; those of the second pair remain widely separated, and are, at about two-thirds of their length, angularly inflected and directed backward. On the side of the body, somewhat behind the middle of the interstice between the second and third pairs of legs, a long bristle is inserted, and underneath, near the third pair of coxæ, stands, as in the male, a straight, lancet-like spine, which is usual in this genus. A little behind the middle of the ventral surface there is a semicircular transverse trabecula in front of the genital orifice; its branches are continued backward and united with the coxal circles of the fourth pair. The mandibles are as in the male.

In the nymph the abdomen terminates in a conical process, which by a narrow fissure is separated into two lobes, each lobe bearing on its outer edge a lanceolate spine and a long bristle on its top.

The hexapod larva is very small, and shows, though in a rudimentary state, the anal appendices of the adult.

The eggs are long, comparatively large, fusiform, transparent, and cribrate.

NOTE.—Several years have elapsed since the publication of this memoir was commenced in December 1886. In the meanwhile I have become better acquainted with the Acarid-fauna of Europe and some other countries. The clearer insight into the leading features of the geographical distribution in general which I have thus acquired, and of which I have given a *résumé* in the "Introduction," has made me sceptical with regard to the validity of some of the species described in this work. As an excuse for the synonymical errors into which I may have fallen in some instances, I may be allowed to plead the many difficulties which the study of Acarids offers to the naturalist in a tropical country, in consequence of the extreme delicacy of the soft-bodied species and their great liability to rapid changes of form and colour. Moreover, owing to the unfortunate circumstance that I only brought

with me to Europe drawings and more or less fragmentary notes of many of the soft-bodied species, I had no opportunity of comparing the types themselves with European forms and the descriptions of authors.

These remarks will perhaps incite future inquirers to fix their attention more particularly on these fragile forms.

The list which follows contains the names of all the species of Acari hitherto described from Mexico and Central America. In it I have incorporated certain corrections and criticisms, which will be found under the names of the species to which they severally refer.

List of the Species of Acariden hitherto described from Mexico and Central America.

Suborder I. ACARINA-TRACHEATA.

Fam. TROMBIDIDÆ, C. L. Koch.

TROMBIDIUM, Latr.

1. Trombidium mexicanum.

Trombidium mexicanum, Stoll, antch, p. 1, Tab. I. figg. 1-1 d.

Hab. MEXICO, Presidio.

This undoubtedly bears a remarkable affinity to *T. dubrueilli*, A. Dugès; yet it cannot be considered to coincide with that species, as it seems to differ not only in size, but also in some minor details as regards the disposition of the hairs and the form of the palpi and mandibles.

2. Trombidium dubrueilli.

Trombidium dubrueilli, Alf. Dugès, La Naturellez, vii. p. 306, t. 8. figg. 1-10.

Hab. MEXICO, Guanajuato, Tupaturo.

3. Trombidium hispidum.

Trombidium hispidum, Stoll, antch, p. 2, Tab. II. figg. 1-1 d.

Hab. GUATEMALA, Retalhuleu.

4. Trombidium nasutum.

Trombidium nasutum, Stoll, antch, p. 2, Tab. III. figg. 1-1 g.

Hab. GUATEMALA, Retalhuleu.

5. *Trombidium quinque-maculatum*.*Trombidium quinque-maculatum*, Stoll, antea, p. 3, Tab. IV. figg. 1-1 c.*Hab.* GUATEMALA, near the city.6. *Trombidium muricola*.*Trombidium muricola*, Stoll, antea, p. 5, Tab. II. figg. 3-3 d.*Trombidium guayacicola*, Stoll, antea, p. 4, Tab. II. figg. 2-2 c.*Hab.* GUATEMALA, Retalhuleu, Antigua.

T. guayacicola and *T. muricola* are, I believe, only colour-varieties of one and the same species. I was led into error by the different habitats, and by the discrepancies between my original drawings, which were made, at a year's interval, at different places, Retalhuleu and Antigua. I propose to drop the name of *T. guayacicola* and to keep only that of *T. muricola*.

7. *Trombidium trilineatum*.*Trombidium trilineatum*, Stoll, antea, p. 4, Tab. I. figg. 2-2 c.*Hab.* GUATEMALA, Antigua.8. *Trombidium albicolle*.*Trombidium albicolle*, Stoll, antea, p. 5, Tab. I. figg. 3, 3 a.*Hab.* GUATEMALA, Antigua.

RHYNCHOLOPHUS, Dugès.

1. *Rhyncholophus erinaceus*.*Rhyncholophus erinaceus*, Stoll, antea, p. 6, Tab. IV. figg. 2-2 d.*Hab.* GUATEMALA, Antigua.

Fam. ACTINEDIDÆ.

ACTINEDA, C. L. Koch.

1. *Actineda baccarum*.*Acarus baccarum*, Linn. Syst. Nat. 10th edit. p. 617.*Acarus vitis*, Schrank, Enum. Ins. Austr. indig. p. 519.*Trombidium cornigerum*, Herm. Mém. Aptérolog. p. 38, t. 2. fig. 9.*Actineda flavicola*, Stoll, antea, p. 7, Tab. V. figg. 1, 1 a.*Actineda antiyuensis*, Stoll, antea, p. 7, Tab. V. figg. 2-2 d.*Actineda retalteca*, Stoll, antea, p. 7, Tab. V. figg. 3-3 c.*Hab.* EUROPE.—GUATEMALA, Retalhuleu, Antigua.—SOUTH AMERICA.

After having studied more attentively the European *A. baccharum*, L. (= *cornigerum*, Herm.), I believe that the above-named Guatemalan forms are but varieties of colour and age of this species, as I cannot find any tangible differences between my original drawings and European specimens, except those of size and colour, which in *Actineda* are of no value. This view is confirmed by the circumstance that Berlese has identified preserved specimens of *Actineda* from Rio Apa (Paraguay) and from Buenos Ayres with the European species, with the remark "parum ab *Europæa diversa*"*. It therefore seems that *A. baccharum*, L. (= *A. vitis*, Schrank, = *Tromb. cornigerum*, Hermann), is one of those fundamental and characteristic types which occupy an extremely extensive geographical area, a fact I have discussed at length in the "Introduction" to the present memoir.

Fam. TETRANYCHIDÆ, Kramer.

TETRANYCHUS, Dufour.

1. *Tetranychus guatemalæ-novæ*.

Tetranychus guatemalæ-novæ, Stoll, antch. p. 8, Tab. VI. figg. 1-1 c.

Hab. GUATEMALA, near the city.

2. *Tetranychus dugesi*.

Tetranychus dugesi, D. Cano y Alencio, La Naturaleza, vii. p. 197, t. 2. figg. 1-5.

Hab. MEXICO. On *Medicago denticulata*, Willd.

Fam. HYDRACHNIDÆ, Neuman.

The corrections in the synonymy of the species of this family described by me are made on the competent authority of Herr Koenike of Bremen, who, at my request, has favoured me with his opinion on them.

ATAX, Fabr.

1. *Atax alticola*.

Atax alticola, Stoll, antch. p. 9, Tab. VII. figg. 1-1 g.

Atax septem-maculatus, Stoll, antch. p. 9, Tab. VIII. figg. 1-1 c.

* A. Berlese (Acar. Austro-Americani, p. 11) identifies his American specimens with *Acarus vitis*, Schrank, which he considers to be synonymous with *Trombidium cornigerum*, Hermann. But as Schrank himself, in his description of *A. vitis* (Zoon. Ins. Austr. indig. p. 519, no. 1067), says that it differs from *A. baccharum*, L., only by the disposition of the hairs on the legs, I think that we must regard *A. vitis*, Schrank, and *Acarus baccharum*, L., as belonging to one and the same species.

Atax septem-maculatus, var. *ypsilon*, Stoll, anteh, p. 10, Tab. IX. figg. 1-1 c.

Hab. GUATEMALA, near the city.

According to Herr Koenike, *A. septem-maculatus* and its var. *ypsilon* are probably nymphal stages of *A. alticola*.

2. *Atax dentipalpis*.

Atax dentipalpis, Stoll, anteh, p. 10, Tab. X. figg. 1-1 d.

Hab. GUATEMALA, near the city.

This, according to the same authority, should be referred to the widely distributed *A. crassipes*, O. F. Müll.

NESÆA, Neuman.

1. *Nesæa alzatei*.

Atax alzatei, A. Dugès, La Naturelle, vi. p. 344, t. 8. figg. 1-19.

Hab. MEXICO, Guanajuato.

This species has been ascribed by its author to the genus *Atax*.

2. *Nesæa guatemalensis*.

Nesæa guatemalensis, Stoll, anteh, p. 11, Tab. X. figg. 2-2 b (♀), and Tab. XI. figg. 1-1 f (♂).

Hab. GUATEMALA, near the city.

3. *Nesæa numulus*.

Nesæa numulus, Stoll, anteh, p. 12, Tab. XI. figg. 2-2 c.

Hab. GUATEMALA, near the city.

LIMNESIA, Neuman.

1. *Limnesia guatemalteca*.

Limnesia guatemalteca, Stoll, anteh, p. 13, Tab. VII. figg. 2-2 e.

Hab. GUATEMALA, near the city.

From the form of the palpi, Herr Koenike considers this to be a nymph, probably of the following species.

2. *Limnesia longipalpis*.

Limnesia longipalpis, Stoll, anteh, p. 13, Tab. IX. figg. 2-2 c.

Hab. GUATEMALA, near the city.

3. *Limnesia puteorum*.*Limnesia puteorum*, Stoll, anteh, p. 14, Tab. VII. figg. 3-3 c.*Hab.* GUATEMALA, Antigua.4. *Limnesia leta*.*Limnesia leta*, Stoll, anteh, p. 14, Tab. VIII. figg. 2-2 d.*Hab.* GUATEMALA, near the city.

Fam. BDELLIDÆ.

BDELLIA, Latr.

1. *Bdella splendida*.*Bdella splendida*, Stoll, anteh, p. 15, Tab. III. figg. 2-2 c.*Hab.* GUATEMALA, near the city.

Fam. EUPODIDÆ.

SCYPHIUS, Koch.

1. *Scyphius maniacus*.*Scyphius maniacus*, Stoll, anteh, p. 17, Tab. VI. figg. 2-2 d.*Hab.* GUATEMALA, Retalhuleu.

Fam. IXODIDÆ.

ARGAS, Latr.

1. *Argas talaje*.*Argas talaje*, Guérin-Ménéville, Revue et Mag. Zool. 1849, p. 342, t. 9.*Hab.* GUATEMALA, Casas Viejas de Guastatoya.2. *Argas turicata*.*Argas turicata*, A. Dugès, La Naturelle, vi. p. 195, t. 4 a. figg. a 1-8.*Hab.* MEXICO, Guanajuato.

A parasite on hogs.

3. *Argas megnini*.*Argas megnini*, A. Dugès, La Naturelle, vi. p. 197, t. 4 a. figg. a 1-8.*Hab.* MEXICO, Guanajuato.

This species is stated to attach itself to the skin in the ears of horses, cattle, and even man.

ORNITHODOROS, Koch.

1. *Ornithodoros coriaceus*.

Ornithodoros coriaceus, C. L. Koch, Arch. f. Naturg. x. Bd. 1, p. 219, no. 1; Übers. d. Arachniden-syst. Heft iv. pp. 12, 31, t. 1. figg. 2, 3.

Hab. MEXICO (*teste Koch*).

Berlese, in his 'Acari Austro-Americani,' p. 23, mentions *Ornithodoros coriaceus*, Koch, from Rio Apa (Paraguay), but without entering into details.

IXODES, C. L. Koch.

1. *Ixodes boarum*.

Ixodes boarum, Stoll, antea, p. 18, Tab. XIII. figg. 1-1 c, and Tab. XIV. fig. 4.

Hab. GUATEMALA, Retalhuleu.

2. *Ixodes pygmaeus*.

Ixodes pygmaeus, C. L. Koch, Arch. f. Naturg. x. Bd. 1, p. 233, no. 18; Übers. d. Arachniden-syst. Heft iv. pp. 22, 107, and t. 22. figg. 81 a, b (♀).

Hab. MEXICO.—BRAZIL (*teste C. L. Koch*).

AMBLIOMMA, C. L. Koch.

1. *Amblyomma mixtum*.

Amblyomma mixtum, C. L. Koch, Arch. f. Naturg. x. Bd. 1, p. 227, no. 17; Übers. d. Arachniden-syst. Heft iv. pp. 74, 75, t. 13. figg. 47 (♂), 48 (♀); Stoll, antea, p. 19, Tab. XII. figg. 1-1 i (♀), 2-2 b (♂).

Hab. MEXICO (*teste C. L. Koch*); GUATEMALA, Retalhuleu, Antigua; NICARAGUA, Chontales; COSTA RICA, Caché.

2. *Amblyomma dissimile*.

Amblyomma dissimile, C. L. Koch, Arch. f. Naturg. x. Bd. 1, p. 225, no. 10 (♂ ♀); Übers. d. Arachniden-syst. Heft iv. pp. 17, 64, 66, t. 12. figg. 37 (♂), 38 (♀).

Hab. MEXICO (*teste Koch*).

3. *Amblyomma tenellum*.

Amblyomma tenellum, C. L. Koch, Arch. f. Naturg. x. Bd. 1, p. 227, no. 16 (♂); Übers. d. Arachniden-syst. Heft iv. pp. 17, 78, 79, t. 14. fig. 51 (♂).

Hab. MEXICO (*teste Koch*).

4. *Amblyomma ovale*.

Amblyomma ovale, C. L. Koch, Arch. f. Naturg. x. Bd. 1, p. 227, no. 20; Übers. d. Arachniden-syst. Heft iv. pp. 18, 79, 80, t. 14. fig. 52 (♂).

Hab. MEXICO (*teste Koch*).

BIOG. CENTR.-AMER., Arachn. Acar., January 1893.

H*

5. **Amblyomma foreli.**

Amblyomma foreli, Stoll, antea, p. 21, Tab. XII. figg. 3-3 δ , and Tab. XIV. figg. 3-3 d (♀).

Hab. GUATEMALA, Retalhuleu.

6. **Amblyomma crassipunctatum.**

Amblyomma crassipunctatum, Stoll, antea, p. 22, Tab. XIV. figg. 1-1 δ (d).

Hab. NICARAGUA, Chontales.

7. **Amblyomma sabanerae.**

Amblyomma sabanerae, Stoll, antea, p. 23, Tab. XIV. figg. 2-2 i (♀).

Hab. GUATEMALA, near Retalhuleu.

Fam. ORIBATIDÆ.

Subfam. *PTEROGASTERINÆ*.

ORIBATA, Latr.

1. **Oribata centro-americana.**

Oribata centro-americana, Stoll, antea, p. 24, Tab. XV. figg. 1-1 f .

Hab. BRITISH HONDURAS, R. Hondo, R. Sarstoon, Belize; GUATEMALA, Antigua, Guatemala city; PANAMA, Volcan de Chiriqui.

2. **Oribata rugifrons.**

Oribata rugifrons, Stoll, antea, p. 25, Tab. XV. figg. 2-2 d (7 nymph, Tab. XV. figg. 3-3 d).

Hab. BRITISH HONDURAS, Belize; GUATEMALA, Retalhuleu.

Subfam. *APTEROGASTERINÆ*.

HOPLOPHORA, C. L. Koch.

1. **Hoplophora retalteca.**

Hoplophora retalteca, Stoll, antea, p. 27, Tab. XV. figg. 4-4 f .

Hab. GUATEMALA, Retalhuleu.

Fam. NICOLETIELLIDÆ.

NICOLETIELLA, R. Canestrini.

1. **Nicoletiella neotropica.**

Nicoletiella neotropica, Stoll, antea, p. 27, Tab. XVI. figg. 1-1 c .

Hab. GUATEMALA, Retalhuleu.

Fam. GAMASIDÆ.

Subfam. UROPODINÆ.

UROPODA, Latr.

1. *Uropoda moneta*.

Uropoda moneta, Walck. & Gervais, Hist. nat. des Ins. Aptères, iii. p. 221, t. 34. fig. 5.

Hab. MEXICO. Found as a parasite on *Polydesmus mexicanus*.

2. *Uropoda echinata*.

Uropoda echinata, Stoll, anteh, p. 28, Tab. XVI. figg. 2-2 c.

Hab. GUATEMALA, Antigua.

3. *Uropoda inæquipunctata*.

Uropoda inæquipunctata, Stoll, anteh, p. 29, Tab. XVI. figg. 3-3 d.

Hab. GUATEMALA, Retalhuleu.

4. *Uropoda discus*.

Uropoda discus, Stoll, anteh, p. 29, Tab. XVII. figg. 4-4 c.

Hab. GUATEMALA, Retalhuleu.

5. *Uropoda centro-americana*.

Uropoda centro-americana, Stoll, anteh, p. 30, Tab. XVII. figg. 1-1 f, and 2-2 b (nymph).

Hab. NICARAGUA, Chontales.

6. *Uropoda piriformis*.

Uropoda piriformis, Stoll, anteh, p. 31, Tab. XVII. figg. 3-3 d.

Hab. MEXICO, Jalapa.

Subfam. GAMASINÆ.

MEGISTHANUS, T. Thorell.

1. *Megisthanus gigantodes*.

Megisthanus gigantodes, Stoll, anteh, p. 31, Tab. XVIII. figg. 1-1 d (♀), 2-2 g (♂).

Hab. GUATEMALA, Aceituno, Guatemala city, Zapote, Retalhuleu, Chelhuiz.

2. *Megisthanus armiger*.

Megisthanus armiger, A. Berlese, Bull. Soc. Ent. Ital. xx. p. 204, t. 9, iv. fig. 1; Stoll, anteh, p. 34,

Tab. XIX. figg. 1-1 e (♂).

Hab. MEXICO, Jalapa.—PARAGUAY, Rio Apa.

CELÆNOPSIS, Berlese.

1. *Celænopsis uropodoides*.

Celænopsis uropodoides, Stoll, anteh., p. 35, Tab. XVI. figg. 4-4 *d*, Tab. XIX. figg. 3-3 *♂* (♀).

Hab. BRITISH HONDURAS, R. Hondo.

2. *Celænopsis megisthanoides*.

Celænopsis megisthanoides, Stoll, anteh., p. 36, Tab. XIX. fig. 4, Tab. XX. figg. 1-1 *♂* (♀)

Hab. PANAMA, Bugaba.

PACHYLÆLAPS, Berlese.

1. *Pachylælaps hæros*.

Pachylælaps hæros, var. *mexicanus*, Stoll, anteh., pp. 37, 38, Tab. XIX. figg. 2-2 *e*.

Pachylælaps hæros, Berlese, Bull. Soc. Ent. Ital. xx. p. 196, t. 8. fig. 2.

Hab. MEXICO, San Andres Tuxtla. [Typus : Matto Grosso, Brazil.]

HOLOSTASPIS, Kolenati.

1. *Holostaspis marginatus*.

Holostaspis marginatus (Hermann), Stoll, anteh., p. 39, Tab. XX. figg. 2-2 *d*.

Hab. GUATEMALA, Retalhuleu; NICARAGUA, Chontales.—SOUTH AMERICA, Brazil, La Plata, Paraguay.

Suborder H. ACARINA-ATRACHEATA.

Fam. SARCOPTIDÆ.

Subfam. ANALGESINÆ.

MEGNINIA, Berlese.

1. *Megninia pteroglossorum*.

Megninia pteroglossorum, Stoll, anteh., p. 40, Tab. XXI. figg. 5-5 *♂*.

Hab. GUATEMALA, Retalhuleu.

PTEROLICHUS, Robin et Mégnin.

1. *Pterolichus momotorum*.

Pterolichus momotorum, Stoll, anteh., p. 41, Tab. XXI. figg. 1-1 *♂* (*♂*), 2, 2 *a* (♀).

Hab. GUATEMALA, Retalhuleu.

PROCTOPHYLLODES, Robin et Ménézin.

1. *Proctophyllodes sialiarum*.

Proctophyllodes sialiarum, Stoll, antea, p. 42, Tab. XXI. figg. 3 (♂), 4-6 c (♀).

Hab. GUATEMALA, Retalhulen.

N.B.—I have noticed the presence of a few other genera in Guatemala, but my notes and drawings respecting them are too fragmentary to allow specific determination. These genera are: *Linopodes*, C. L. Koch, *Gamasus*, Latr. (sensu stricto), *Pteroptus*, Duf., *Tyroglyphus*, Latr.

INDEX.

[Names in small capitals refer to Families &c.; those in roman type to the chief reference to each species included in the work; those in italics to species incidentally mentioned, synonym, &c.]


	Page		Page		Page
ACARINA-TRACHEATA	39, 52	Atax septem-maculatus,		Limacina longipalpis	13, 47
ACARINA-TRACHEATA	1, 44	var. ypsilon	10	— puterum	14, 48
Acerus laevigatus	46	— septem-maculatus,		Limacodes	53
— marginatus	39	var. ypsilon	47		
— nitidus	45, 46			Megisthanus armiger	34
ACTINEDA	7, 45	BOELLA	15, 48	MEGISTHANTS	31, 51
Actineta	45	Boella	16	Megisthanus	33, 54
— angustatus	7	— splendida	15, 48	— armiger	31, 51
— angustatus	45	BOELLIDÆ	15, 48	— gigantodes	31, 51
— buccinum	45			MACONIA	45, 52
— buccinum	45	CERKNOPIA	35, 52	— pteroglossum	45, 52
— curvatum	45	— megisthanoides	35, 52	Marcia	26
— flavula	7	— subvires	36	— trisculata	26
— flavula	45	— uropodoides	35, 52		
— reticulata	7			NEREA	11, 47
— reticulata	45	DERMANYSINÆ	30	Nereis	12
— vitis	45, 46	Diplogynium	35	— albatel	47
ACTINERIDÆ	6, 45			— guatemalensis	11, 47
AMALYOMMA	10, 46	EUPODIÆ	17, 48	— numulus	12, 47
Amalyomma	18, 21			Nicotia	27
— crassipunctatum	22, 50	GAMARINÆ	28, 51	NICOLETIELLA	27, 50
— dissimile	49	GAMARINÆ	31, 51	— cornuta	27
— fovei	21, 50	Gammarus	30, 53	— neotropica	27, 50
— mixtum	19, 49	— calceptratum	30	NICOLETIELLIDÆ	27, 50
— mixtum	21, 22	— fuscum	30		
— ovalis	49			ORBATA	24, 50
— sabulosus	33, 50	HOLOSTAPHIS	30, 52	— centro-americana	24, 50
— senilis	49	— marginatus	30, 52	— rufifrons	25, 50
ANALOGINÆ	40, 52	— marginatus, var. americanus	30	— rufifrons	26
APTEROGASTERINÆ	37, 50	HOPLOPHORA	27, 50	Orbata	24
ARGAS	48	— nitens	27	ORIBATIDÆ	24, 50
— megini	46	— reticulata	27, 50	ORIBATOPODO	40
— talaja	48	— reticulata	26	— coriacas	40
— turicata	48	HYDRACHNIDÆ	9, 46		
ATAX	9, 41			PACHYLELAPS	37, 52
Atax	10, 47	IXODES	18, 49	— laevis	37, 52
— albicollis	9, 41	— bonum	18, 49	— laevis	38
— albicollis	47	— pygmaeus	49	— laevis, var. mexicanus	37, 38
— alutici	47	IXODES	18, 48	— laevis, var. mexicanus	39
— crassipes	47			PHACOTOPHYLLOIDES	42, 53
— dentipalpis	10, 47	LENXIA	13, 47	— stellarum	42, 53
— septem-maculatus	9	— guatemalensis	12, 47	PTEROGASTERINÆ	24, 50
— septem-maculatus	10, 46, 47	— lata	14, 48	PTEROLICHES	41, 53

INDEX.

55

	Page		Page		Page
<i>Pteridichus biemarginatus</i>	42	TETRANTCHIDÆ	8, 46	<i>Trembidium basatum</i>	2, 44
— <i>monotermum</i>	41, 53	<i>Tetrantchus</i>	8, 46	— <i>quinque-maculatum</i>	5, 45
— <i>monotermum</i>	42	— <i>dupoi</i>	46	— <i>trilineatum</i>	4, 45
— <i>phyllipsectus</i>	42	— <i>guatemalensis-noti</i>	8, 46	TYROGLYPHIDÆ	39
— <i>ramphastius</i>	42	TROMBIDIDÆ	1, 44	<i>Tyroglyphus</i>	39, 53
PTEROPTES	39	TROMBIDUM	1, 44		
<i>Pteroptus</i>	53	<i>Trombidium</i>	10	UROPODA	28, 51
		— <i>albicollis</i>	5, 45	<i>Uropoda</i>	30, 39
RHYNCHOLOPHUS	6, 45	— <i>cornigerum</i>	45, 46	— <i>caudata</i>	30
<i>Rhyncholophus</i>	16, 17	— <i>dohrnii</i>	44	— <i>centro-americana</i>	30, 51
— <i>electoralis</i>	16	— <i>gonysavicola</i>	4	— <i>discus</i>	30, 51
— <i>crinaceus</i>	6, 45	— <i>gonysavicola</i>	45	— <i>echinata</i>	28, 51
		— <i>hippidum</i>	2, 44	— <i>inquinatata</i>	29, 51
SARCOPTIDÆ	39, 52	— <i>mexicanum</i>	1, 44	— <i>moneta</i>	51
<i>Sarcoptides</i>	39	— <i>merinense</i>	2, 5	— <i>piriformis</i>	31, 51
SCYPHIUS	17, 48	— <i>muricola</i>	6, 45	UROPODIDÆ	28, 51
— <i>manicus</i>	17, 48				

JUL 15 1919



1-1a. *TETRANYCHUS MERIANAE*

2-2c. *TETRANYCHUS TRITICI*

3-3a. *TETRANYCHUS ALBIPENNIS*

MU



1a



3



1d



1



3a



2



1b



1c



3b



2a



2c

Fig. 1. Ticks: 1 - *Ixodes ricinus*; 2 - *Ixodes ricinus*; 3 - *Ixodes ricinus*.

Fig. 1a, b, c, d - details of the legs of the ticks.

Fig. 2a, b, c - details of the legs of the ticks.

110

Fig. 1. *Phytolaela* sp.

Fig. 2. *Phytolaela* sp.



1a.



1b.



1



1c.



1d.



1e.



1f.



2a.



2b.



2



2c.



2d.



2e.



2f.



3



3a.





3b.

Fig. 3. *Phytolaela* sp.

Fig. 4. *Phytolaela* sp.

Fig. 5. *Phytolaela* sp.

20



40

100
100
100
100



1a.



1.



1c



2a



2.



2a



2c



2d



2e

70






Fig. 1. *U. (U.)* *...* Fig. 2. *U. (U.)* *...*

70



THE JOURNAL OF THE ENTOMOLOGICAL SOCIETY OF AMERICA

Vol. 1, No. 1, 1910

175





Fig. 1. 187. 1. Mite (Dorsal view). 2. Tick (Dorsal view). 3. Ticks (Dorsal view).

1000




1. ♀ *ACANTHOSOMA DENTIFALX*.

2. ♀ *NESETA GUATEMALENSIS*.

(From the original figures in the original work.)

NU




1d




2






1. 1. 2. 3. 4. 5. 6. 7. 8. 9. 10. 11. 12. 13. 14. 15. 16. 17. 18. 19. 20. 21. 22. 23. 24. 25. 26. 27. 28. 29. 30. 31. 32. 33. 34. 35. 36. 37. 38. 39. 40. 41. 42. 43. 44. 45. 46. 47. 48. 49. 50. 51. 52. 53. 54. 55. 56. 57. 58. 59. 60. 61. 62. 63. 64. 65. 66. 67. 68. 69. 70. 71. 72. 73. 74. 75. 76. 77. 78. 79. 80. 81. 82. 83. 84. 85. 86. 87. 88. 89. 90. 91. 92. 93. 94. 95. 96. 97. 98. 99. 100. 101. 102. 103. 104. 105. 106. 107. 108. 109. 110. 111. 112. 113. 114. 115. 116. 117. 118. 119. 120. 121. 122. 123. 124. 125. 126. 127. 128. 129. 130. 131. 132. 133. 134. 135. 136. 137. 138. 139. 140. 141. 142. 143. 144. 145. 146. 147. 148. 149. 150. 151. 152. 153. 154. 155. 156. 157. 158. 159. 160. 161. 162. 163. 164. 165. 166. 167. 168. 169. 170. 171. 172. 173. 174. 175. 176. 177. 178. 179. 180. 181. 182. 183. 184. 185. 186. 187. 188. 189. 190. 191. 192. 193. 194. 195. 196. 197. 198. 199. 200. 201. 202. 203. 204. 205. 206. 207. 208. 209. 210. 211. 212. 213. 214. 215. 216. 217. 218. 219. 220. 221. 222. 223. 224. 225. 226. 227. 228. 229. 230. 231. 232. 233. 234. 235. 236. 237. 238. 239. 240. 241. 242. 243. 244. 245. 246. 247. 248. 249. 250. 251. 252. 253. 254. 255. 256. 257. 258. 259. 260. 261. 262. 263. 264. 265. 266. 267. 268. 269. 270. 271. 272. 273. 274. 275. 276. 277. 278. 279. 280. 281. 282. 283. 284. 285. 286. 287. 288. 289. 290. 291. 292. 293. 294. 295. 296. 297. 298. 299. 300. 301. 302. 303. 304. 305. 306. 307. 308. 309. 310. 311. 312. 313. 314. 315. 316. 317. 318. 319. 320. 321. 322. 323. 324. 325. 326. 327. 328. 329. 330. 331. 332. 333. 334. 335. 336. 337. 338. 339. 340. 341. 342. 343. 344. 345. 346. 347. 348. 349. 350. 351. 352. 353. 354. 355. 356. 357. 358. 359. 360. 361. 362. 363. 364. 365. 366. 367. 368. 369. 370. 371. 372. 373. 374. 375. 376. 377. 378. 379. 380. 381. 382. 383. 384. 385. 386. 387. 388. 389. 390. 391. 392. 393. 394. 395. 396. 397. 398. 399. 400. 401. 402. 403. 404. 405. 406. 407. 408. 409. 410. 411. 412. 413. 414. 415. 416. 417. 418. 419. 420. 421. 422. 423. 424. 425. 426. 427. 428. 429. 430. 431. 432. 433. 434. 435. 436. 437. 438. 439. 440. 441. 442. 443. 444. 445. 446. 447. 448. 449. 450. 451. 452. 453. 454. 455. 456. 457. 458. 459. 460. 461. 462. 463. 464. 465. 466. 467. 468. 469. 470. 471. 472. 473. 474. 475. 476. 477. 478. 479. 480. 481. 482. 483. 484. 485. 486. 487. 488. 489. 490. 491. 492. 493. 494. 495. 496. 497. 498. 499. 500. 501. 502. 503. 504. 505. 506. 507. 508. 509. 510. 511. 512. 513. 514. 515. 516. 517. 518. 519. 520. 521. 522. 523. 524. 525. 526. 527. 528. 529. 530. 531. 532. 533. 534. 535. 536. 537. 538. 539. 540. 541. 542. 543. 544. 545. 546. 547. 548. 549. 550. 551. 552. 553. 554. 555. 556. 557. 558. 559. 560. 561. 562. 563. 564. 565. 566. 567. 568. 569. 570. 571. 572. 573. 574. 575. 576. 577. 578. 579. 580. 581. 582. 583. 584. 585. 586. 587. 588. 589. 590. 591. 592. 593. 594. 595. 596. 597. 598. 599. 600. 601. 602. 603. 604. 605. 606. 607. 608. 609. 610. 611. 612. 613. 614. 615. 616. 617. 618. 619. 620. 621. 622. 623. 624. 625. 626. 627. 628. 629. 630. 631. 632. 633. 634. 635. 636. 637. 638. 639. 640. 641. 642. 643. 644. 645. 646. 647. 648. 649. 650. 651. 652. 653. 654. 655. 656. 657. 658. 659. 660. 661. 662. 663. 664. 665. 666. 667. 668. 669. 670. 671. 672. 673. 674. 675. 676. 677. 678. 679. 680. 681. 682. 683. 684. 685. 686. 687. 688. 689. 690. 691. 692. 693. 694. 695. 696. 697. 698. 699. 700. 701. 702. 703. 704. 705. 706. 707. 708. 709. 710. 711. 712. 713. 714. 715. 716. 717. 718. 719. 720. 721. 722. 723. 724. 725. 726. 727. 728. 729. 730. 731. 732. 733. 734. 735. 736. 737. 738. 739. 740. 741. 742. 743. 744. 745. 746. 747. 748. 749. 750. 751. 752. 753. 754. 755. 756. 757. 758. 759. 760. 761. 762. 763. 764. 765. 766. 767. 768. 769. 770. 771. 772. 773. 774. 775. 776. 777. 778. 779. 780. 781. 782. 783. 784. 785. 786. 787. 788. 789. 790. 791. 792. 793. 794. 795. 796. 797. 798. 799. 800. 801. 802. 803. 804. 805. 806. 807. 808. 809. 810. 811. 812. 813. 814. 815. 816. 817. 818. 819. 820. 821. 822. 823. 824. 825. 826. 827. 828. 829. 830. 831. 832. 833. 834. 835. 836. 837. 838. 839. 840. 841. 842. 843. 844. 845. 846. 847. 848. 849. 850. 851. 852. 853. 854. 855. 856. 857. 858. 859. 860. 861. 862. 863. 864. 865. 866. 867. 868. 869. 870. 871. 872. 873. 874. 875. 876. 877. 878. 879. 880. 881. 882. 883. 884. 885. 886. 887. 888. 889. 890. 891. 892. 893. 894. 895. 896. 897. 898. 899. 900. 901. 902. 903. 904. 905. 906. 907. 908. 909. 910. 911. 912. 913. 914. 915. 916. 917. 918. 919. 920. 921. 922. 923. 924. 925. 926. 927. 928. 929. 930. 931. 932. 933. 934. 935. 936. 937. 938. 939. 940. 941. 942. 943. 944. 945. 946. 947. 948. 949. 950. 951. 952. 953. 954. 955. 956. 957. 958. 959. 960. 961. 962. 963. 964. 965. 966. 967. 968. 969. 970. 971. 972. 973. 974. 975. 976. 977. 978. 979. 980. 981. 982. 983. 984. 985. 986. 987. 988. 989. 990. 991. 992. 993. 994. 995. 996. 997. 998. 999. 1000.

Digitized by Google




1. I. IXODES RICINUS.



1. *AMPLOMMA CRASSIFUNCTUM*
SABANERA

2. *AMPLOMMA PUBESCENS*
IXTAPALAPA


NU



1-11 CHIRATA CENTIO AMERICANA
EUSIFRONS.

1-4 LAEVA OF CHIRATA ST
HOLIFRONS ESTALTEA


nd



1-10. *BOULETIA A. NEOTROPICA*.
11-14. *BOULETIA A. NEOTROPICA*.

15-18. *BOULETIA A. NEOTROPICA*.
19-22. *BOULETIA A. NEOTROPICA*.


Digitized by Google



1. 4. NERISTHANS NGANTODES.

40

Digitized by Google



11659 CELENOPSIS MECISTRANOIDS
 11660 HOL STAPH. MARGINATUS.

no

